Baština Akademije nauka i umjetnosti Bosne i Hercegovine

Proceedings of the Conference on March 14 - International Day of Mathematics

Vuković, Mirjana, urednik; Nurkanović, Mehmed, urednik

2024-12-26

Academy of Sciences and Arts of Bosnia and Herzegovina

https://bastina.anubih.ba/handle/123456789/798

Preuzeto s Baštine Akademije nauka i umjetnosti Bosne i Hercegovine

https://bastina.anubih.ba/

PROCEEDINGS OF THE CONFERENCE ON MARCH 14 – INTERNATIONAL DAY OF MATHEMATICS
Special Editions ANUBiH, Book CCXVI, OPMN Book 30
Sarajevo 2024, **p. 63–67** DOI: 10.5644/PI2024.216.08

PROPERTIES OF SOME SPECTRA OF SUPERPOSITION OPERATORS

SANELA HALILOVIĆ

ABSTRACT. We consider the nonlinear superposition operator F in Banach spaces of sequences l_p, generated by the function $f(s,u)$. We analyze the Rhodius spectra $\sigma_R(F)$ and the Neuberger spectra $\sigma_N(F)$ of these operators F generated by the function

$$f(s,u) = a(s) + \phi(u),$$

where $(a(s))_{s\in\mathbb{N}}$ is a sequence from l_p $(1 \le p \le \infty)$, and $\phi(u)$ is a continuous function in $\mathbb{R}$. Some connections between the property of the function $\phi(u)$ and the corresponding spectra $\sigma_R(F)$ and $\sigma_N(F)$ are given in this paper. There are also a few examples that verify proposed theorems.

1. INTRODUCTION

In this paper we consider the Rhodius and Neuberger spectra of superposition operators in Banach spaces l_p $(1 \le p \le \infty)$. The superposition operators have an important place in many mathematical problems and also there are various applications in mathematical physics, mathematical economics, mathematical biology and so on. Let $f(s,u)$ be a function defined on $\mathbb{N}\times\mathbb{R}$ with values in $\mathbb{R}$. Given a function $x = x(s)$, by applying f, we get the function $f(s,x(s))$ and this function generates an operator F

$$F(x(s)) = f(s,x(s)). \tag{1.1}$$

This operator (1.1) is called the superposition operator, composition operator or Nemytskii operator. We take $x = x(s)$ a sequence from the l_p spaces of sequences $(1 \le p \le \infty)$, which are the Banach spaces equipped with the standard norm. It is known that the spectrum of a linear operator has many useful properties. For nonlinear operators F, the notion spectrum of F is a wider concept, based on the property of $\lambda I - F$ being a regular map.

For the class of all continuous operators F on a Banach space X, denoted by $\mathfrak{C}(X)$, the following definition of Rhodius spectrum has been introduced.

Definition 1.1. *For the continuous operator $F : X \to X$ the set*

$$\rho_R(F) = \{\lambda \in \mathbb{R} : \lambda I - F \text{ is bijective and } (\lambda I - F)^{-1} \in \mathfrak{C}(X)\}$$

2020 *Mathematics Subject Classification.* 47J10, 47H30.

Key words and phrases. superposition operator, spectra, nonlinear operator, Rhodius spectrum, Neuberger spectrum.

is called the Rhodius resolvent set and

$$\sigma_R(F) = \mathbb{R} \setminus \rho_R(F)$$

is the Rhodius spectrum.

Thus, a point $\lambda \in \mathbb{R}$ belongs to $\rho_R(F)$ if and only if $\lambda I - F$ is a homeomorphism on X. The Neuberger spectrum of nonlinear operators was proposed for $\mathfrak{C}^1(X)$, the class of continuously Fréchet differentiable operators on Banach space X.

Definition 1.2. *For an operator $F : X \to X$, which is continuously Fréchet differentiable, the Neuberger resolvent set is defined by*

$$\rho_N(F) = \left\{\lambda \in \mathbb{R} : \lambda I - F \text{ is bijective and } (\lambda I - F)^{-1} \in \mathfrak{C}^1(X)\right\}$$

and the set $\sigma_N(F) = \mathbb{R} \setminus \rho_N(F)$ is called the Neuberger spectrum of F.

A point $\lambda \in \mathbb{R}$ belongs to $\rho_N(F)$ if and only if $\lambda I - F$ is a diffeomorphism on X.
Some useful properties of these spectra are:

- If F is a linear operator $\Rightarrow \sigma_R(F) = \sigma_N(F) = \sigma(F)$,
- If $F0 = 0 \Rightarrow \sigma_R(F) \subseteq \sigma_N(F)$,
- If the underlying space is complex, then $\sigma_N(F)$ is nonempty.

More information about various nonlinear spectral theories can be found in [1]. The conditions of acting, continuity and differentiability of the superposition operator in Banach spaces l_p, are given in the following three theorems from [2].

Theorem 1.1. *Let $1 \le p, q < \infty$. Then the following properties are equivalent:*

- *the operator F acts from l_p to l_q;*
- *there are functions $a(s) \in l_q$ and constants $\delta > 0, n \in \mathbb{N}, b \ge 0$, for which*

$$|f(s,u)| \le a(s) + b|u|^{\frac{p}{q}} \ (s \ge n, |u| < \delta);$$

- *for any $\epsilon > 0$ there exists a function $a_\epsilon \in l_q$ and constants $\delta_\epsilon, > 0, n_\epsilon \in \mathbb{N}$, $b_\epsilon \ge 0$, for which $\|a_\epsilon(s)\|_q < \epsilon$ and*

$$|f(s,u)| \le a_\epsilon(s) + b_\epsilon|u|^{\frac{p}{q}} \ (s \ge n_\epsilon, |u| < \delta_\epsilon).$$

Theorem 1.2. *Let $1 \le p, q < \infty$ and let the superposition operator* (1.1)*, generated by the function $f(s,u)$, acting from l_p to l_q. Then this operator is continuous if and only if each of the functions is continuous for every $s \in \mathbb{N}$.*

Theorem 1.3. *Let $1 \le p, q < \infty$ and the operator F generated by the function $f(s,u)$ acting from l_p into l_q. The operator F is differentiable at $x_0 \in l_p$ if and only if $f'_u(s,\cdot)$ is continuous at x_0 for almost all $s \in \mathbb{N}$.*

2. Main Results

The Rhodius and Neuberger spectra are defined for a continuous operator F, that is, $F \in \mathfrak{C}(l_p)$, so the function $\phi(u)$ has to be continuous function on $\mathbb{R}$.

Theorem 2.1. *Let the superposition operator $F : l_p \to l_p$ be generated by the function $f(s,u) = a(s) + \phi(u)$, where $(a(s))_s$ is a sequence from the space l_p $(1 \le p \le \infty)$. If $\phi(u)$ is a bijective function, then $0 \in \rho_R(F)$ and $0 \notin \sigma_R(F)$*

Proof. For $\lambda = 0$ the operator $\lambda I - F$ becomes $-F$. If $\phi(u)$ is a bijection then the functions $f(s,u) = a(s) + \phi(u)$ and $-f(s,u) = -a(s) - \phi(u)$ are bijective for every $s \in \mathbb{N}$. It follows that the operator F, generated by f and the operator $-F$, generated by $-f$ are bijective. Since $-f$ is a bijective function, it follows that its inverse $(-f)^{-1}$ exists and it is also a bijective function. The function $-f(s,u)$ is a bijective and continuous for every s and hence $(-f)^{-1}(s,u)$ is a bijective and continuous function for every s. From the Theorem 1.2 we conclude the operator $(-F)^{-1}$ generated by $(-f)^{-1}$ is a continuous operator. So, we show that operator $-F$ is bijective and $(-F)^{-1} \in \mathfrak{C}(l_p)$. It means that $0 \in \rho_R(F)$ and $0 \notin \sigma_R(F)$. □

The Neuberger spectrum is defined for $F \in \mathfrak{C}^1(l_p)$, so the function $\phi(u)$ has to be continuously differentiable on $\mathbb{R}$. Then the derivative $f'_u(s,u) = \phi'(u)$ is a continuous function for all $s \in \mathbb{N}$ and from Theorem 1.3 it follows the operator F is continuously Fréchet differentiable.

Theorem 2.2. *Let the superposition operator $F : l_p \to l_p$ be generated by the function $f(s,u) = a(s) + \phi(u)$, where $(a(s))_s$ is a sequence from the space l_p $(1 \le p \le \infty)$. Let $\phi(u)$ be a bijective and continuously differentiable function.*
a) If $\phi'(u) \neq 0, \forall u$, then $0 \in \rho_N(F)$ and $0 \notin \sigma_N(F)$.
b) If there exists u_0 such that $\phi'(u_0) = 0$, then $0 \notin \rho_N(F)$ and $0 \in \sigma_N(F)$.

Proof. a) The function $\phi(u)$ is a continuously differentiable function, so $\phi'(u)$ exists for every $u \in \mathbb{R}$ and it is a continuous function. Since the function ϕ is bijective, its inverse ϕ^{-1} exists and it is also bijective. If $\phi'(u) \neq 0, \forall u$, then in virtue of the inverse function theorem (see [6]), the ϕ^{-1} is a differentiable function and

$$(\phi^{-1})'(y) = \frac{1}{\phi'(u)} \tag{2.1}$$

holds, where $\phi(u) = y$ and $\phi^{-1}(y) = u$.
We have $f'_u(s,u) = (a(s) + \phi(u))'_u = \phi'(u) \neq 0$ and

$$(f^{-1})'_u = \frac{1}{f'_u(s,u)} = \frac{1}{\phi'_u(u)}. \tag{2.2}$$

This derivative $(f^{-1})'_u$ is continuous in u because $\phi'(u)$ is a continuous function and $\phi'(u) \neq 0$ for every u. That is why, according to the Theorem 1.3, $(f^{-1})'_u$ generates a continuous operator $(F^{-1})'$, i.e. $(F^{-1}) \in \mathfrak{C}^1(l_p)$. Then clearly, $((-f)^{-1})'_u$ is also a continuous function and $(-F)^{-1} \in \mathfrak{C}^1(l_p)$. In the proof of Theorem 2.1 we have already shown that $-F$ is a bijective operator and now we see that $(-F)^{-1} \in \mathfrak{C}^1(l_p)$, so we have proved that $0 \in \rho_N(F)$ and $0 \notin \sigma_N(F)$.
b) If there exists u_0 such that $\phi'(u_0) = 0$, then the function ϕ^{-1} is not differentiable at $y_0 = \phi(u_0)$ and consequently, the partial derivatives $(f^{-1})'_u$ and $(-f^{-1})'_u$ are not continuously differentiable functions. Hence, $(-F)^{-1} \notin \mathfrak{C}^1(l_p)$ and this means that $0 \notin \rho_N(F)$ and $0 \in \sigma_N(F)$. □

2.1. **Examples**

Here we give two examples of nonlinear superposition operators and their Rhodius and Neuberger spectra, which illustrate and verify these theorems (Theorem 2.1 and Theorem 2.2).

Example 2.1. *Let the operator $F : l_p \to l_p$ be generated by the function*

$$f(s,u) = a(s) + u^n, \tag{2.3}$$

where $n \geq 3$ and n is an odd number. This function $\phi(u) = u^n$ is bijective. In [4] it was found that $\sigma_R(F) = (0,\infty)$, so $0 \notin \sigma_R(F)$. So, in this example we see that the function $\phi(u)$ is bijective and $0 \notin \sigma_R(F)$ and this confirms Theorem 2.1.
The Fréchet derivative of the operator F generated by (2.3)*, at $x_0 = (x_1, x_2, \ldots)$ along $h = (h_1, h_2, \ldots)$ is:*

$$F'(x_0)h = (nx_1^{n-1}h_1, nx_1^{n-1}h_2, \ldots).$$

In [3] we found $\sigma_N(F) = [0,\infty)$, so $0 \in \sigma_N(F)$. Here, the function $\phi(u) = u^n$ is continuously differentiable ($\phi'(u) = n \cdot u^{n-1}$), but its inverse $\phi^{-1}(u) = \sqrt[n]{u}$ is not differentiable at $u = 0$ $\left((\phi^{-1})'(u) = \frac{1}{n} \cdot \frac{1}{\sqrt[n]{u^{n-1}}}\right)$. In this case the function $\phi(u) = u^n$ is bijective and $\phi'(u) = n \cdot u^{n-1}$. Therefore, there exists $u_0 = 0$, such that $\phi'(u_0) = \phi'(0) = 0$. In virtue of the Theorem 2.2 it follows that $0 \in \sigma_N(F)$. Hence, in this example Theorem 2.2 is verified.

Example 2.2. *Let the operator $F : l_p \to l_p$ be generated by the function*

$$f(s,u) = a(s) + \sqrt[n]{u}, \tag{2.4}$$

where $n \geq 3$ and n is an odd number. This function $\phi(u) = \sqrt[n]{u}$ is bijective and the function $f(s,u)$ is bijective for every $s \in \mathbb{N}$, so the operator F is bijective. In [4] we found that $\sigma_R(F) = (0,\infty)$, so $0 \notin \sigma_R(F)$. Hence,in this example the function $\phi(u) = \sqrt[n]{u}$, (n-odd number) is bijective and $0 \notin \sigma_R(F)$ and it agrees with Theorem 2.1.
The Fréchet derivative of the operator F generated by (2.4)*, at $x_0 = (x_1, x_2, \ldots)$ along $h = (h_1, h_2, \ldots)$ is:*

$$F'(x_0)h = \left(\frac{1}{n\sqrt[n]{x_1^{n-1}}}h_1, \frac{1}{n\sqrt[n]{x_2^{n-1}}}h_2, \ldots\right).$$

The function $-f(s,u)$ is bijective and the operator $-F$ is bijective. We have

$$-f(s,u) = -a(s) - \sqrt[n]{u}$$

and

$$(-f)^{-1}(s,u) = (-a(s) - u)^n.$$

Let us find the partial derivative of $(-f)^{-1}$ with respect to the variable u :

$$((-f)^{-1})'_u = n \cdot (-u - a(s))^{n-1} \cdot (-u - a(s))'_u = -n \cdot (-u - a(s))^{n-1}. \tag{2.5}$$

The function (2.5) *is continuous for all $s \in \mathbb{N}$ and from Theorem 1.3 it follows that the operator $(-F)^{-1}$ is continuously differentiable. Here we see that $-F$ is a bijective operator and $(-F)^{-1} \in \mathfrak{C}(l_p)$. Now from Definition 1.2 it follows $0 \in \rho_N(F)$ and $0 \notin$*

$\sigma_N(F)$. In this case we have the function $\phi(u) = \sqrt[n]{u}$ which is bijective and

$$\phi'(u) = (\sqrt[n]{u})'_u = \frac{1}{n} \cdot \frac{1}{\sqrt[n]{u^{n-1}}} \neq 0, \forall u \in \mathbb{R}. \tag{2.6}$$

Therefore, from Theorem 2.2 it follows that $0 \in \rho_N(F)$ and $0 \notin \sigma_N(F)$. So, in this example we also get the validation of Theorem 2.2.

3. CONCLUSION

In this paper we observe the class of superposition operators $F : l_p \to l_p$, generated by the function $f(s,u) = a(s) + \phi(u)$. We find out how the fact that the function $\phi(u)$ is bijective affects the Rhodius and Neuberger spectra of the operator F, generated by the function $f(s,u)$. We conclude that it affects the corresponding spectra in regards to whether $\sigma_R(F)$ and $\sigma_N(F)$ contain 0. In [5] we found that if the function $\phi(u)$ is not a bijection, then $0 \in \sigma_R(F)$ and $0 \in \sigma_N(F)$. Our further goal is to investigate how some other properties of these spectra of nonlinear superposition operators, such as closedness, boundedness etc, depend on the properties of their generating function $f(s,u)$.

REFERENCES

[1] J. Appell, E. De Pascale and A. Vignoli, *Nonlinear Spectral Theory*, Walter de Gruyter, 2004.
[2] F. Dedagić and P. P. Zabreiko, *On the superposition operator in l_p spaces*, Sibir. Mat. Zhurn., vol.28, No.1, (1987), pp. 86-98.
[3] S. Halilović and R. Vugdalić, *The Neuberger Spectra of Nonlinear Superposition Operators in the Spaces of Sequences*, Journal of the International Mathematical Virtual Institute, vol.4, (2014), pp. 97-119.
[4] S. Halilović and R. Vugdalić, *The Rhodius Spectra of Some Nonlinear Superposition Operators in the Spaces of Sequences*, Adv. Math., Sci.J., vol.3, no. 2, (2014), pp. 83-96.
[5] S. Halilović, *Some Spectra of Superposition Operators Generated by an Exponential Function*, Communications in Mathematics and Applications, vol.12, no. 1 (2021), pp. 221-229.
[6] W. Rudin, *Principles of Matemathematical Analysis*, Mc Graw-Hill, 1976.

(Received: May 18, 2024)
(Revised: July 27, 2024)

Sanela Halilović
University of Tuzla
Faculty of Natural Sciences and Mathematics
Urfeta Vejzagića 4
75000 Tuzla
Bosnia and Herzegovina
e-mail: *sanela.halilovic@untz.ba*