Baština Akademije nauka i umjetnosti Bosne i Hercegovine

Cardiovascular diseases and oral health – the impact of pregnant women’s oral health on children’s cardiovascular health

Mesihović Dinarević, Senka

2022-06-03

Academy of Sciences and Arts of Bosnia and Herzegovina

https://bastina.anubih.ba/items/7352c280-9809-4d6f-97ed-551ca13e1944

Preuzeto s Baštine Akademije nauka i umjetnosti Bosne i Hercegovine

https://bastina.anubih.ba/

Chapter VII
PROJECT PUBLICATIONS

Journal of Cardiovascular Diseases & Diagnosis

Mesihović-Dinarević, J Cardiovasc Dis Diagn 2017, 5:3
DOI: 10.4172/2329-9517.1000268

Opinion Article OMICS International

Prevention of Cardiovascular Disease from an Early Age

Senka Mesihović-Dinarević*
Policlinic Eurofarm, Sarajevo, Bosnia and Herzegovina

Introduction

Atherosclerosis is the most frequent disease of arteries characterized by lumen reduction of blood vessel due to local thickening of internal blood vessel by plaque/atheroma [1-3]. It is now one of the leading causes of death in developed countries. Of the ten most common causes of death, diseases of the heart and blood vessels, as a result of atherosclerosis, accounts for six of them. Atherosclerosis begins in childhood, the patient goes a long time without developing symptoms, increasing with age and at about 50 years of age, atherosclerosis seriously begins to threaten health. As a cardiovascular disease, atherosclerosis is an interdisciplinary problem that is treated by: cardiologists, neurologists, epidemiologists, nutritionists etc.

Pathophysiology

Pathophysiology of this disease concerns damaged cells that line the inner surface of the arteries the endothelium, due to chemical or mechanical damage [4]. Elevated blood cholesterol, smoking, or elevated homocysteine are examples of chemical and mechanical damage to the cells as well as high blood pressure, damage of the catheter during diagnostic procedures or even infections. Hypertension causes the formation of plaque in places where electricity hits the wall of the blood vessel and creates vortices, namely: the aortic arch, bifurcation, arteries, initial parts of the heart or coronary arteries, but also long-term or frequent spasms and immunological mechanisms lead to endothelium damage. The defect site develops inflammation and damaged endothelial cells secrete a variety of substances that attract other cells in the environment, cholesterol, platelets, smooth muscle cells, and by various interactions do accumulate in the walls of arteries. The first stage of the development of atherosclerosis occurs in the form of a "fatty streak." It is totally reversible, which means that with cessation of activities associated with harmful pathogens, endothelial cells can recover completely. However, if exposure to adverse events continues, atherosclerotic plaque grows further narrowing lumen. Consequently, reduced blood flow and tissue oxygen supply become insufficient. In the blood vessel, itself due to reduced elasticity, an increase of blood pressure occurs [5]. Elevated blood pressure can lead to: bursting in blood of the court or its stratification with the formation of aneurysms or plaque, which can calcify and slim down the wall of the vessel with the ability of forming a clot. For atherosclerotic disease, itself plaque composition rather than its size is much more important. Plaque is primarily built of lipids, has a thin cap, hat plaque, and is easier to shoot. If thrombi are formed they can clot arteries. In contrast, larger plaque with a hard cap, which contains less fat, can rarely explode, because it is more stable.

Important risk factors

The most important risk factors for the development of atherosclerotic disease are: hyperlipidaemia, hypertension, smoking, diabetes, high fibrinogen, male sex at a younger and middle age, menopause in women taking oral contraceptives or hormone replacement therapy only with the presence of other risk factors, excessive weight, increased levels of homocysteine, physical inactivity, heredity and immune response in some diseases [6,7].

Consequences of atherosclerosis

The Consequences of atherosclerosis are: coronary or ischemic heart disease, especially myocardial infarction, cerebrovascular disease and cerebrovascular accidents (80% of all heart attacks and stroke due to atherosclerosis), the narrowing or blockage of peripheral arteries, carotid arteries, particularly the legs, which can even lead to the development of gangrene [8].

The process of hypertension starts in childhood. Etiopathogenetically it is multifactorial, a possible course and repercussions for health are longstanding and irreversible [9]. Normotension offers important data in contribution to health and increased blood pressure in childhood represents the site for preventive paediatric action. In a study which included 500 children, a correlation of birth weight and blood pressure in the paediatric population was conducted at the Paediatric clinic of Clinical University Centre of Sarajevo, during April-June 2003 [10]. The conclusion was that blood pressure measurement in children could serve for the detection of cardiovascular disease precursors' ages 8 to 9. The correlation of birth weight and systolic pressure exists, but not significantly. The main factors that influence birth weight are: sex (boys were heavier than girls up to 98 g $p<0.05$), gestational age (preterm were lighter for 600 g $p<0.001$), and smoking of pregnant mothers (newborns were for 219 g of lower birth weight in relation to mothers who did not smoke $p<0.01$). Obese participants (important hypertension factor) have increased systolic blood pressure for 5.38 mmHg in relation to those participants with normal blood tension ($p<0.01$).

Dislipoproteinaemias and repercussions on the myocardium and blood vessels in the paediatric population, represent an imperative for modern investigation. Basic investigation of lipoproteins, apolipoprotein's metabolism, the biology of the atheromatosus process development in cells, and the role of genetics in the development of disease of coronary arteries is necessary in the field of preventive cardiology. Arteriosclerosis, multifactorial in its etiopathogenesis, course and repercussions, demands action from a team of physicians with the aim of early detection and treatment of dislipoproteinemias and the reduction of the development of risk factors for coronary disease. As atherosclerosis is one of the leading causes of death in society, it represents a responsibility for all paediatricians who should do a screening of lipid levels for all children aged two with a positive family history, as well as children in schools. Dislipoproteinemia is detectable in neonatal age, so the data from Sarajevo's study from 1991, one of the first investigations in the field of preventive cardiology in Europe, points to an elevation of total cholesterol levels, as well as an increase of apo B lipoprotein fraction [11]. In this study, the lipoprotein serum

***Corresponding author:** Senka Mesihović-Dinarević, Policlinic Eurofarm, Sarajevo, Bosnia and Herzegovina, E-mail: dsenka@bih.net.ba

Received March 27, 2017; **Accepted** April 17, 2017; **Published** April 21, 2017

Citation: Mesihović-Dinarevi S (2017) Prevention of Cardiovascular Disease from an Early Age. J Cardiovasc Dis Diagn 5: 268. doi: 10.4172/2329-9517.1000268

Copyright: © 2017 Mesihović-Dinarević S. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Mesihović-Dinarevi S (2017) Prevention of Cardiovascular Disease from an Early Age. J Cardiovasc Dis Diagn 5: 268. doi: 10.4172/2329-9517.1000268

level was evaluated in 163 patients without positive family history of cardiovascular risk factors. The control group was comprised of 32 healthy children. In children with cardiovascular disease (congenital heart disease, rhythm disorders, rheumatic fevers), in relation to the control group, an increase of c lipid fraction (triglyceride and LDLP) with a decrease of protective fraction of HDL with statistical significance was proven, adding to increased risk for cardiovascular diseases.

Since there is no specific cure for atherosclerosis, the best way to avoid this disease is prevention. In the United States and Western Europe, where risk factors have been preventively eliminated for a long time, mortality rates "fell" below 50%. Prevention included avoiding risk factors: instead of obesity subjects should have normal body weight according to sex, age and height, work out or quick walk at least half an hour each day, because 40% of our bodies are comprised of skeletal muscles that are designed to move [6]. The primary intention of prevention is to preclude the occurrence of risk factors for atherosclerosis, and the secondary is to prevent the development or aggravation of the illness along with the reduction or control of existing risks. Primary prevention should begin as early as possible, even in childhood, creating a healthy diet, eliminating smoking, regular physical activity, which will prevent or at least slow the development of atherosclerosis. In modern medicine, there are a growing number of studies that show that children are overweight, which is certainly an adolescent risk factor for many chronic diseases including: cardiovascular, diabetes type 2, orthopaedic and psychological illnesses. The epidemic of obesity is one of the most serious health problems of today. During the past two decades, the prevalence of obesity in European countries has increased threefold. Fifty percent of adults today are overweight and a third of the European population is obese. Statistics from the Centre for Disease Control and Prevention /CDC/ also speak to the tripling of the number of obese people in the last 20 years [12]. Sixteen percent of children and adolescents aged 6-19 years are overweight. Approximately 60% to 85% of obese children become obese adults, leading to an earlier and more frequent occurrence of chronic non-communicable disease. Though genetic and hormonal factors are possible causes of children being overweight and obese, excessive food intake and low physical activity are undoubtedly the main reasons for the emergence of obesity. Sitting in front of TV and computer whilst consuming calorie rich fast foods and sweetened beverages create long-term imbalances between the introduction and consumption of energy in the body. The result of this imbalance is being overweight [13,14]. Parameters for the most precise estimate of obesity is the body mass index (BMI), which is the ratio of body weight and the square of body height in kg/m^2. Obese people have a BMI greater than the 95^{th} percentile. Factors related to the start of obesity are: a modified diet, reduced physical activity, and increased inactivity. The alarming trend of the expanding epidemic of obesity, particularly an increase in prevalence among young people, presents each community with a problem that has enormous economic and social consequences [15-20]. Global measures for the prevention of obesity for the countries of the European region are given in the European Charter of countering Obesity in 2006 [21].

Finding the most effective preventive measures for obesity in each country requires accurate epidemiological data on the number of obese children and adolescents, and their dietary habits and activity, which was one of the goals important research carried out in 2008-2010 in Canton Sarajevo [14]. The results of this massive study of risk factors for cardiovascular disease (the largest of it's kind in Bosnia and Herzegovina), which was carried out in Canton Sarajevo, from the earliest ages-children in kindergartens, primary and secondary school students and the working age population, practically ages 0-65, a total of 42,828 respondents were surveyed. Through results from this research, we got the data on: risk factors for cardiovascular disease and atherosclerosis, including the level of physical activity in school environments, and the possibility of prevention, diagnosing and treating to this problem using a modern approach through cognitive behavioural aspects. For the assessment of the nutritional state of children and adolescents we used the body mass index, the degree of nutrition is obtained automatically based on CDC criteria: BMI<5 percentile malnutrition. BMI: 5 and 85 percentile normal BW, BMI: 85-95 over nutrition, BMI greater than 95 percentile obesity. The representative sample of students in elementary and secondary schools in Canton Sarajevo were selected *via* random sample selection. The number of respondents in all elementary (1-8) and secondary (1-4) grades was balanced. Pupils were interviewed in written form. The survey forms were originally designed and included questions about their habits in food intake (frequency, quantity and types) and fluids, and the frequency and intensity of physical activity. The survey forms for primary and secondary schools contained similar questions but were adapted to the age group of specific subjects: lower elementary grades, higher grades of elementary school and secondary school students. Measurements of anthropometric parameters: height and weight, was done for all subjects. Height was measured using a vertical scale in centimetres (cm), and the results are rounded to 0.5 cm. Body weight was measured by electronic floor scales in kilograms (kg), and the results are rounded to 0.5 kg. The research team consisted of two physicians and two graduate nurses. The research was concluded by 2 teams. Subjects voluntarily participated in the survey and measurements the data after being input into the information system was made anonymous from all subjects. The survey and measurements of anthropometric parameters was performed in a total of 3608 students from Canton Sarajevo of which: 2329 were from 9 elementary schools and 1279 from 6 high schools. The results were: about 1/4 of children in the first 4 years are obese, and in higher classes 1/5 of pupils. According to the results of the questioner a majority children do not eat healthy food at home. There is a problem at school concerning eating habits: pupils from lower classes eat food from school and a majority eat food from bakeries (43.31%). Physical activity was documented in 19.92% of younger children. In the older groups of pupils there is a greater percentage of pupils who sit in front the TV and PC for up to 2 hours (30.56%) they were also sitting during learning which means the majority of the day. A great number of pupils (58.15%) eats sweets every day between meals (Tables 1-6).

Conclusion

In Conclusion, the question raised is: what to do about the prevention of cardiovascular disease? Why do schools need to worry about health? As a society, we value good health. Good health is necessary for effective learning. Healthy students become healthy, productive citizens. Schools are the places where we spend most of our youth so we have to initiate a change. The teamwork of an environmental society and new policies are needed. As a priority, we should create a place for physical activity and nutrition in schools; develop funded prevention programs and a systematic approach to the problem. Point to the problem of obesity in an adequate manner, and create partnerships with value to the social community in creating a healthy lifestyle. Schools can provide: quality physical education classes, healthy nutrition campaigns, parenting education, psychosocial education/intervention with nutrition strategy. Prevention is the key! It is necessary to establish dialogue in cardiovascular medicine. Prevention of obesity consists of nutrition conducted according to modern guidelines in relation to the input of: carbohydrates, fats, proteins, vitamins and fluids. It is necessary to

Citation: Mesihović-Dinarevi S (2017) Prevention of Cardiovascular Disease from an Early Age. J Cardiovasc Dis Diagn 5: 268. doi: 10.4172/2329-9517.1000268

School class	Total number of students	M	F	BMI classification			
				Under Weight (%)	Normal weight (%)	Over weight (%)	Obesity (%)
I-IV Elementary School	1077	546	531	T: 20,86	T: 55,26	T: 12,28	T: 11,58
				M: 7,72	M: 28,68	M: 7,47	M: 6,77
				F: 13,14	F: 26,58	F: 4,81	F: 4,81
V-VIII Elementary School	1252	680	572	T: 9,16	T: 69,80	T: 13,07	T: 8,00
				M: 4,65	M: 37,98	M: 7,49	M: 3,69
				F: 4,51	F: 31,82	F: 5,58	Ž: 4,31
I-IV High School	1279	531	748	T: 6,76	T: 80,43	T: 9,55	T: 3,24
				M: 1,31	M: 34,15	M: 5,18	M: 1,47
				F: 5,45	F: 46,28	F: 4,37	F: 1,77
All schools	3608	1757	1851	T: 12,49	T: 68,74	T: 11,86	T: 6,86
				M: 4,63	M: 33,60	M: 6,71	M: 1,32
				F: 7,86	F: 35,14	F: 5,15	F: 5,54

Table 1: BMI classification of students according to grades, schools and gender.

School and class	Homemade sandwich (%)	Sandwich bought at school (%)	Food from the bakery (%)	Snacks (%)	'I dont eat at school' (%)
I-IV Elementary School	30,69	44,63	5,04	17,68	1,98
V-VIII Elementary School	16,57	24,62	42,67	10,96	5,27
I-IV High School	2,70	41,09	42,01	8,89	5,37
All schools	16,65	36,78	29,90	12,57	4,20

Table 2: Distribution of the quality nutrition of students in school.

School class	Water (%)	Juices (%)	Milk (%)	Sodas (%)
I-IV Elementary School	50,88	23,19	23,37	2,6
V-VIII Elementary School	54,94	21,71	12,45	11,87
I-IV High School	49,35	16,77	13,19	20,72
All schools	51,82	20,65	16,43	11,05

Table 3: Distribution of types of beverages that students drink during the day.

School class	Candys every day (%)	Sometimes and rarely (%)
I-IV Elementary School	30,89	59,11
V-VIII Elementary School	64,53	33,47
I-IV High School	80,85	19,15
All schools	58,15	31,24

Table 4: Distribution of prevalence of candy consuming.

School class	Every day activity (%)	Activity only on sport class (%)	Activity 2-3x per week (%)	Rarely (%)
I-IV Elementary School	19,92	39,05	40,36	0,00
V-VIII Elementary School	36,42	27,56	24,52	11,50
I-IV High School	46,89	29,19	18,08	5,65
All schools	31,07	31,93	27,65	8,51

Table 5: Distribution of the degree of sports activities.

School class	< 1 hour (%)	1-2 hours (%)	2-3 hours (%)	3 hours (%)
I-IV Elementary School	23,60	36,62	18,51	20,89
V-VIII Elementary School	13,81	30,56	27,74	27,96
I-IV High School	18,70	33,59	23,12	24,42

Table 6: Distribution of the length of time spent in front of computer and TV.

maintain regular physical activity, both in school and in free time. It is also necessary to promote of continuous education about healthy eating and a healthy lifestyle (activity, not smoking) in all primary and secondary schools in the country, the engagement of teams of family doctors to monitor the nutritional status of school children and youth.

References

1. Berenson GS, Blonde CV, Faris RP (1979) Cardiovascular disease risk factor variable during the first year of life. Am J Dis Child 133: 1049-57.
2. Berenson GS, Srinivasan SR, Hunter SM, Nicklas TA, Freedman DS, et al. (1989) Risk factors in early life as predictors of adult heart disease: The Bogalusa heart study. Am J Med Sci 298: 141-151.
3. Geer JC, McGill, He Strong JP (1961) The fine structure of human atherosclerotic lesions. Am J Pathol 38: 263-269.
4. Dinarević S (1994) The pathogenesis of atherosclerosis-a review. Brit J Cardiol 67: 241-246.
5. Moore S (1985) Pathogenesis of atherosclerosis. Metabolism 34: 13-16.
6. Dhuper S, Buddhe S, Patel S (2013) Managing cardiovascular risk in overweight children and adolescents. Paediatr Drugs 15: 181-190.
7. Berenson GS, Srinivasan SR, Bao W, Newman WP III, Tracy RE, et al. (1998) Association between multiple cardiovascular risk factors and atherosclerosis in children and young adults. The Bogalusa Heart Study. N Engl J Med 338: 1650-1656.
8. Newman WP, Freedman DS, Voors AW, Gard PD, Srinivasan SR, et al. (1986) Relation of serum lipoprotein levels and systolic blood pressure to early atherosclerosis. The Bogalusa Heart Study. New Eng J Med 314: 138-144.
9. National High Blood Pressure Education Program Working Group on High Blood Pressure in Children and Adolescents (2004) The fourth report on the diagnosis, evaluation, and treatment of high blood pressure in children and adolescents. Paediatrics 114: 555-76.
10. Dinarević S, Mulaosmanović V (2005) Primary prevention of Hypertension in Sarajevo Children: Role of Adiposity. 29th UMEMPS Congress Union of Middle Eastern and Mediterranean Paediatric Societies, pp. 154-156.
11. Dinarević S, Mesihović H, Simeunović S, Zulić I (1994) Dyslipoproteinaemia in Children with Heart Disease. Intercontinental Cardiol 3: 126-9.
12. Centers for Disease Control and Prevention (2015) Over weight and Obesity: Data & Statistics. Accessed on: April 3, 2017.
13. Kapur G, Ahmed M, Pan C, Mitsnefes M, Chiang M, et al. (2010) Secondary hypertension in overweight and stage 1 hypertensive children: A Midwest Paediatric Nephrology Consortium report. J Clin Hypertens 12: 34-39.
14. Dinarević S, Hasanbegović S (2010) Problem of obesity in children and youth in Canton Sarajevo. Pediatr Res 68: 1091.
15. WHO Europe (2007) The challenge of obesity in the WHO European Region and the strategies for response. Copenhagen: WHO Regional Office for Europe.

Citation: Mesihović-Dinarevi S (2017) Prevention of Cardiovascular Disease from an Early Age. J Cardiovasc Dis Diagn 5: 268. doi: 10.4172/2329-9517.1000268

16. Daniels SR, Arnett DK, Eckel RH, Gidding SS, Hayman LL, et al. (2005) Overweight in children and adolescents: Pathophysiology, consequences, prevention and treatment. Circulation 111: 1999-2012.

17. Berenson GS, Wittingly WA, Tracy RE, Newman WP, Srinivasan SR, et al. (1992) Atherosclerosis of the aorta and coronary arteries and cardiovascular risk factors in persons aged 6 to 30 years and studied at necropsy (The Bogalusa Heart Study). Am J Cardiol 70: 851-858.


18. McNiece KL, Gupta-Malhotra M, Samuels J, Bell C, Garcia K, et al. (2007) National High Blood Pressure Education Program Working Group: Left ventricular hypertrophy in hypertensive adolescents: Analysis of risk by 2004 National High Blood Pressure Education Program Working Group staging criteria. Hypertension 50: 392-5.

19. Torrance B, McGuire KA, Lewanczuk R, McGavock J (2007) Overweight, physical activity and high blood pressure in children: a review of the literature. Vasc Health Risk Manag 3: 139-149.

20. Genovesi S, Antolini L, Giussani M, Federico P, Sara G, et al. (2008) Usefulness of waist circumference for the identification of childhood hypertension. J Hypertens 26: 1563-70.

21. http://www.euro.who.int/document/E89567.pdf 15–17 November 2006.

Citation: Mesihović-Dinarevi S (2017) Prevention of Cardiovascular Disease from an Early Age. J Cardiovasc Dis Diagn 5: 268. doi: 10.4172/2329-9517.1000268


Innovinc
DEDICATED TO YOUR SUCCESS
Challenges in Cardiology:
Cardiology Congress 2018
August 16-18, 2018 | Rome, Italy
https://cardiology.innovincconferences.com/
Venue: Hotel Rome
Pisana Via della Pisana,
374 00163 Roma RM, Italy

Challenges in Cardiology:
Cardiology Congress 2018
August 16-18, 2018 | Rome, Italy

Innovinc
DEDICATED TO YOUR SUCCESS

The challenges of detecting risk factors for the development of atherosclerosis

Senka Mesihović-Dinarević, *Lutvo Sporišević, Berislav Topić, Sanja Jurišić, Senad Saric, Britt Gritt, Vjekoslav Krželj, Anes Jogunčić, Samir Prohić and Aida Ramić*

Policlinic Eurofarm, Sarajevo, Committee for Cardiovascular Pathology, Academy of Sciences and Arts Bosnia and Herzegovina

The most frequent disease of the arteries is atherosclerosis which is characterized by lumen reduction of blood vessels due to local thickening of internal blood vessels caused by plaque / atheroma [1-3]. Atherosclerosis is now one of the leading causes of death in developed countries. The most important risk factors for the development of atherosclerotic disease are: hyperlipidaemia, hypertension, smoking, diabetes, high fibrinogen, excessive weight and physical inactivity [4,5]. Some authors pointed out the possible connection between parodontal disease in pregnant women with risk of preterm delivery, newborns of low gestational age with low birth masses and possible cardiovascular diasese [6-10]. Bearing this in mind, there is ongoing study in the Balkan region with the aim of investigating more predictors of early cardiovascular risk /increased body mass index, high values of blood pressure and thickening of intima-media carotids complex/ in comparison to children whose mothers had good oral health during pregnancy. Preliminary study data: mean age of 43 pregnant women is 30.7+/-5.7 years, 90.3% pregnancy runs properly, KEP index:12.32+/-5.7, plak index 0.312, restored teeth 65.62%; eating habits: 48.4% dairy products, fruit 64.5%, vegetables 22.6%, meat 41.9%, fish in 35.2%. Regular dental therapy can decrease frequency of caries, peridontal disease in pregnant women, the frequency of prematurity, low birth weight (with all it's potential complications, decreasing financial costs of neonatal intensive care management and cardiovascular repercussions on newborn's health). A cardiovascular-oral health data base for the Balkan region can be used as a geographic, demographic and epidemiologic source of information for the detection and identification of new potential risk factors of individuals for preterm delivery and possible atherosclerosis. Primary prevention of atherosclerosis should begin as early as possible, during pregnancy, in childhood, creating a healthy way of life, which will be able to prevent or at least slow the development of atherosclerosis.

References

1. Berenson GS, Blonde CV, Faris RP: Cardiovascular disease risk factor variable during the first year of life. American Journal of Disease in Children, 1979; 133:1049-1057.
2. Berenson GS, Srinivasan SR, Mac D et al: Risk factors in early life as predictors of adult heart disease: The Bogalusa heart study, Am.J.Med.Sci.1989; 298:141-151.
3. Geer JC, McGill, He Strong JP: The fine structure of human atherosclerotic lesions, An. J. Pathol. 1961; 38:263-9.
4. Dhuper S, Buddhe S, Patel S. Managing Cardiovascular Risk in Overweight Children and Adolescents. Paediatric Drugs, 2013, 15(3):181-90. doi: 10.1007/s40272-013-0011-y.
5. Berenson GS, Srinivasan SR, Bao, Newman WP, III, Tracy RE, Wattigney WA. Association between multiple cardiovascular risk factors and atherosclerosis in children and young adults. The Bogalusa Heart Study. N Engl J Med 1998; 338:1650–6.
6. Offenbacher S, Katz V, Fertik G, Collins J, Boyd D, Myynor G at all. Periodontal infection as a possible risk factor of preterm low birth weight . J Periodontal 1996;67:1103-1113.

7. Soroey M, Avanbadejo P, Savage K, Oluwole A. Association between periodontal disease and pregnancy outcomes. Odontostomatol. Trop. 2015; 38 (152):5-16.
8. Turton M, Africa CW. Further evidence for periodontal disease as a risk indicator for adverse pregnancy outcomes. Int Dent J. 2016 Dec 17. doi: 10.1111/idj.12274.
9. Trivedi S, Lal N, Singhal R. Periodontal disease during pregnancy. J Orofacial Sci 2015; 7:67.
10. Govindaraju P, Venugopal S, Shivakumar MA, Sethuraman S, Ramaiah SK, Mukundan S. Maternal periodontal disease and preterm birth: A case-control study. J Indian Soc Periodontol. 2015;19(5):512-515. doi: 10.4103/0972-124X.164751

Biography

Senka Mesihović-Dinarević was born in 1958-Sarajevo.Faculty of Medicine Sarajevo 1982. 1982-2016 worked at the Paediatric Clinic-Clinical University Centre Sarajevo. MSc 1985, paediatrician age 30, subspecialty in paediatric cardiology: Sarajevo, Belgrade, London. PhD 1991, Professor of Paediatrics 2006. 1994. Member of AEPC, 1995. BPCA.1995-2011, lecturer: London MRCP: 2003-2016 Director of Paediatric Clinic. 2000 FESC, 2008 Member of Academy of Sciences and Arts of Bosnia and Herzegovina, 2009. Chairman of the Committee of Cardiovascular pathology, Honorary Doctorate of Letters, Cambridge England 2014. 2016. Member of the European Academy of Science and Arts; over 477 papers, cited 218 times in the ICI Web of Science.

Vol.1 No.1 Septembar 2018. Udruženje kardiologa u Bosni i Hercegovini (Knjiga Abstrakta) pp. 1-178

Udruženje kardiologa u Bosni i Hercegovini
The Association of Cadiologists of B&H

8. KONGRES UDRUŽENJA KARDIOLOGA U BiH



Knjiga Abstrakta

ISSN

Knjiga Abstrakta

Izdavač:
Udruženje kardiologa u Bosni i Hercegovini
71000 Sarajevo
Bosna i Hercegovina

Glavni urednik:

Prof. dr sc. Mirza Dilić,
Redovni profesor, Medicinski fakultet Univerziteta u Sarajevu

Recenzenti:

Prof. dr sc. Zumreta Kušljugić,
Redovni profesor, Medicinski fakultet Univerziteta u Tuzli

Prof. dr sc. Šekib Sokolović
Redovni profesor, Medicinski fakultet Univerziteta u Sarajevu

Copyright©2018. Udruženje kardiologa u Bosni i Hercegovini

Dizajn i štampa:
Mirbergraf d.o.o. Sarajevo

Tiraž:
300

Nacionalna i univerzitetska biblioteka, Bosne i Hercegovine, Sarajevo

ISSN 2566-4719 (print)

Knjiga Abstrakta; Udruženje kardiologa u Bosni i Hercegovini

Knjiga Abstrakta

ORAL AND CARDIOVASCULAR HEALTH- THE CHALLENGES OF DETECTING RISK FACTORS FOR THE DEVELOPMENT OF ATHEROSCLEROSIS

Senka Mesihović-Dinarević*, Lutvo Sporišević, Berislav Topić, Sanja Jurišić, Senad Saric, Grit Kirsten-Sarić, Vjekoslav Krželj, Anes Jogunčić, Samir Prohić, Aida Ramić

Policlinic Eurofarm, Sarajevo, Committee for Cardiovascular Pathology, Academy of Sciences and Arts, Sarajevo, Bosnia and Herzegovina

The most frequent disease of the arteries is atherosclerosis which is characterized by lumen reduction of blood vessels due to local thickening of internal blood vessels caused by plaque/atheroma /1-3. / As a cardiovascular disease, atherosclerosis is an interdisciplinary problem that is treated by: cardiologists, neurologists, epidemiologists, nutritionists etc. Atherosclerosis is now one of the leading causes of death in developed countries. It begins in childhood, goes a long time without manifestings symptoms, increasing with age it begins to seriously threaten health. The most dangerous risk factors for the development of atherosclerotic disease are: hyperlipidaemia, hypertension, smoking, diabetes, high fibrinogen, excessive weight and physical inactivity /4,5/. Some authors pointed out the possible connection between parodontal disease in pregnant women with the risk of preterm delivery, newborns of low gestational age with low birth masses and possible cardiovascular diasese. Bearing this in mind, there is now an ongoing study in the Balkan region with the aim of investigating more prominent predictors of early cardiovascular risk/increased body mass index, high values of blood pressure and the thickening of the intima-media carotids complex in comparison to children whose mothers had good oral health during pregnancy /6-10/. Regular dental therapy/care can decrease the frequency of the occurance of caries and peridontal disease in pregnant women, the frequency of prematurity and low birth weight (with all it's potential complications, decreasing the financial costs of neonatal intensive care management and the cardiovascular repercussions on newborn's health). A cardiovascular-oral health data base for the Balkan region can be used as a geographic, demographic and epidemiologic source of information for the detection and identification of new potential risk factors of individuals for preterm delivery and possible atherosclerosis. Primary prevention of atherosclerosis should begin as early as possible, during pregnancy, in childhood, creating a healthy way of life, which will be able to prevent or at least slow the development of atherosclerosis.

Journal of Cardiovascular Diseases & Diagnosis

Dinarevic et al. J Cardiovasc Dis Diagn 2018, 6:6
DOI: 10.4172/2329-9517.1000342

Research Article Open Access

The Challenges of Detecting Risk Factors for the Development of Atherosclerosis

Dinarević SM*, Topić B, Jurišić S, Prohić S, Sporišević L, Saric S, Sarić GK, Krželj V, Jogunčić A and Ramić A

Committee for Cardiovascular Pathology, Academy of Sciences and Arts, Sarajevo, Bosnia and Herzegovina

Abstract

The most frequent disease of the arteries is atherosclerosis which is characterized by lumen reduction of blood vessels due to local thickening of internal blood vessels caused by plaque/atheroma. As a cardiovascular disease, atherosclerosis is an interdisciplinary problem and one of the leading causes of death in developed countries. It begins in childhood, goes a long time without manifesting symptoms, increasing with age it begins to seriously threaten health. The most dangerous risk factors for the development of atherosclerotic disease are: Hyperlipidaemia, hypertension, smoking, diabetes, high fibrinogen, excessive weight and physical inactivity.

Keywords: Atherosclerosis; Hyperlipidaemia; Hypertension; Myocardial infarction; Parodontal disease

Introduction

The primary intention of prevention of atherosclerosis is to preclude the occurrence of risk factors for atherosclerosis, and the secondary is to prevent the development or aggravation of the illness along with the reduction or control of existing risks. Primary prevention should begin as early as possible, even in childhood, creating a healthy diet, eliminating smoking, regular physical activity, which will prevent or at least slow the development of atherosclerosis. The consequences of atherosclerosis are: coronary or ischemic heart disease, especially myocardial infarction, cerebrovascular disease and cerebrovascular accident (80% of all heart attacks and brain due to atherosclerosis), narrowing or blockage of peripheral arteries, carotid arteries. Since there is no specific cure for atherosclerosis, the best way to prevent this disease, is prevention.

Some authors pointed out the possible connection between parodontal disease in pregnant women with the risk of preterm delivery, new-borns of low gestational age with low birth masses and possible cardiovascular disease [1-7].

Oral health

Appropriate oral health reflects and influences general health and the quality of life. The oral cavity is an integral part of the human organism and therefore there exists a great connection between oral health and systemic health. Not only do some systemic diseases such as diabetes, osteoporosis, HIV infection, trisomy 21 have a predisposition for periodontitis but an opposite applies. Susceptibility to certain systemic disease is higher in patients with periodontitis than in healthy people: chronic periodontitis is a risk factor for future cardiovascular disease, pregnant women with chronic periodontitis have more frequent have preterm birth and new-borns have a low birth weight. The explanation for the pathophysiological mechanisms of paradont focus and systemic disease is associated with elevated levels of circulating pro-inflammatory cytokines and prostaglandins derived from: diseased parodont, gram negative bacteria and their endotoxin-like substances, that appear from subgingival biofilms immediately entering the bloodstream. The dominant problems of everyday dental practice are: caries, periodontal disease, occlusal abnormalities, the relationship of oral and general health and a holistic approach to the patient. Caries and periodontal disease are of an infectious aetiology therefore the prevention of dental caries and periodontal disease means preventing odontogenic focuses. A periodontal pocket is a risk factor for the development or worsening of systemic - focal disease, because the infection is always present in it; a pocket flora is various, massive, virulent and penetrates the soft wall of the pocket, the pocket is under constant mechanical stimulation during chewing, swallowing and speech, all of which favour the penetration of bacteria into circulation and the formation of transient bacteremia. Dental caries and periodontal disease are the most common and significant oral disease, they can cause and aggravate numerous other disease: of the cardiovascular system (infective endocarditis, atherosclerosis, myocarditis and myocardial infarction), of the respiratory system (pneumonia, chronic obstructive pulmonary disease, bronchial asthma and pulmonary abscess), neurological disorders (cerebral infarction and cerebral abscess), diabetes mellitus, rheumatoid arthritis, Alzheimer's disease, and other illnesses) [8]. Interdisciplinary cooperation in the elimination of potential negative effects of periodontal infections will result in better systemic health Atherosclerosis is the basis for all cardiovascular disease. Periodontal pathogens can directly infect the vascular endothelium and atherosclerotic plaque causing inflammation. Then, they are capable of producing a variety of virulence factors (adhesions, haemolysis), which have adverse effects on the vascular system resulting in platelet aggregation and adhesion; in addition, lipid clusters are formed with deposits of cholesterol that contribute to atheroma formation. The treatment of chronic periodontitis reduces systemic inflammation factors.

Pregnancy

Pregnancy is a state in which there are complex physical and physiological changes, which have important effects on multiple systems of organs. High levels of circulating oestrogen during pregnancy are associated with high incidence of gingivitis and gingival hyperplasia or certain forms of periodontal disease [9]. It is believed that approximately 40% of pregnant women have a certain form of periodontal disease [8-10]. Offenbacher et al. first suggested a possible link between periodontal disease and risks for child delivery of a low

***Corresponding author:** Dinarević SM, Committee for Cardiovascular Pathology, Academy of Sciences and Arts, Sarajevo, Bosnia and Herzegovina, Tel: +387 33 560-700; E-mail: dsenka@bih.net.ba

Received October 01, 2018; **Accepted** October 25, 2018; **Published** October 28, 2018

Citation: Dinarević SM, Topić B, Jurišić S, Prohić S, Sporišević L, et al. (2018) The Challenges of Detecting Risk Factors for the Development of Atherosclerosis. J Cardiovasc Dis Diagn 6: 342. doi: 10.4172/2329-9517.1000342

Copyright: © 2018 Dinarević SM, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Dinarević SM, Topić B, Jurišić S, Prohić S, Sporišević L, et al. (2018) The Challenges of Detecting Risk Factors for the Development of Atherosclerosis. J Cardiovasc Dis Diagn 6: 342. doi: 10.4172/2329-9517.1000342

gestational age or small birth weight [11]. Many researchers suggest a possible link between periodontal disease in pregnant women with a risk for preterm delivery, respectively, the birth of babies with low birth weight. Researches show that in pregnant women with periodontal disease there is a 2-7 times higher risk for prematurity. Morre et al. found a large number of foetal deaths among mothers with periodontal disease [12]. However, a focal infection may affect prematurity (birth before 37 weeks gestation) of neonates and neonatal reduction in weight (weight <2500 gm). Bacteraemia of periodontal pathogens trasplacentary can lead to intrauterine infection. Proinflammatory cytokines release the (LPS) endotoxin that precipitates in premature labour. LPS - bacterial lipopolysaccharides, are the major molecular component of the outer membrane of Gram-negative bacteria and serve as a physical barrier providing the bacteria protection from its surroundings. That is why dentists need to motivate, educate and instruct pregnant women towards a higher level of oral hygiene and to repair all dento-oral lesions in dental therapeutic procedures, especially periodontal pockets, thereby reducing the number of premature births. It is believed that the Gram-negative anaerobic bacteria, present in the periodontal tissue, can be a source for endotoxin and lipopolysaccharides, that lead to high levels of inflammatory mediators - interleukin- 6 (IL-6), interleukin 8 (IL-8), Interleukin - 1 beta (IL-1β), and prostaglandin E2 (PGE2) and tumor necrosis factor-alpha (TNF-α), which are transferred to the uterus, cervix and placenta causing premature birth or the birth of children of a small birth weight. Jeffcoat et al. whilst investigating the connection between periodontal disease of pregnant women and preterm birth in group of 1313 pregnant women, found that a moderate to severe form of periodontal disease, diagnosed in early pregnancy, was associated with an increased risk for premature delivery, independently of other traditional risk factors for prematurity [13]. During pregnancy, there is an increased susceptibility to caries due to: the increased acidity of the oral cavity, the increased consumption of refined sugars and poor oral hygiene [14]. Caries bacteria in children are usually transmitted by direct transmission through the mother's saliva. Mothers with high titres of Streptococcus mutants in their saliva are going to substantially transmit the bacteria to their baby – by vertical transmission, creating conditions for early childhood caries. Of course, the time and frequency of the transmission of bacteria, the child's preference for the accumulation of bacteria on its teeth, the composition and flow of the child's saliva, the amount of refined sugar in the baby's food, are all significant predictors of early children's caries [15]. The incidence of births of premature infants and new-borns of a small birth weight is between 5-18%, depending on the geographical area and population characteristics. Due to the immaturity of their organ systems, premature babies and infants of a small birth weight are among in vulnerable group of infants - complications due to prematurity are the leading cause of death in children under five years of age [16]. It is very important to determine risk factors that can lead to the risk of the prematurity of new-borns and new-borns of a small birth weight, i.e. with knowledge of the risk factors, it is possible to substantially eliminate or reduce the risk of premature baby birth or the birth of children with a small weight and to decrease the rate of perinatal mortality and possible complications. Preterm children or infants of a small birth weight exhibit a higher incidence of cardiovascular risk factors (obesity, hypertension, dyslipidaemia), and type 2 diabetes mellitus [16]. Animal and epidemiological studies indicate that conditions of elevated levels of glucocorticoids intrauterine during life, programme the hypothalamus-pituitary-adrenal gland axis that plays a key role in the higher incidence of cardiovascular risk in premature infants and children of a small birth weight [17]. Apart from the role of microbiome mouth (microbiome - all microbes, their genome and mutual interaction in a particular environment) as a risk factor for premature delivery or the birth of new-born of small birth weight, it is possible that the microorganisms of the oral cavity condition chronic inflammation that can represent an atherosclerotic cardiovascular risk factor. Adequate prenatal care should include oral health care of pregnant women, i.e. for pregnant women, there is a need to point out the importance of practicing regular oral hygiene and the need for periodic or as many as or as frequently as needed dental check-ups. The sufficient screening/screening of oral health status of pregnant women is not carried out in daily work, so with screening status of oral health in a greater number of pregnant women, including the assessment of oral hygiene, we would be able to timely identify pregnant women who have dental caries respectively periodontal disease [18]. With timely dental treatment we could reduce the incidence of dental caries and periodontal disease in pregnant women and may reduce the incidence of preterm delivery and the birth of new-borns of a small birth weight, an early childhood caries and predictors of early atherosclerotic cardiovascular risk (increased body mass index, blood pressure and thickening of the carotid intima-media complex). In a cohort of children, who are preterm or have a low birth weight, a certain number of children age 3 have a greater body mass index, a higher value of systolic and diastolic blood pressure, as well as a thickening of the intima-media complex of the carotid artery with incipient signs of cardiovascular system disease [19].

Insufficient insight into the possible pathological implications of the oral health status of pregnant women to premature expression of cardiovascular risk factors in children, initiated this research.

The impact of oral health of pregnant women on the cardiovascular health of children is a Project within the South-eastern European region that runs by the Committee of the Cardiovascular Disease Department of Medical Sciences of Academy of Arts and Sciences of Bosnia and Herzegovina. During 2017, the first phase of research was completed according to plan/lasted for 12 months. In this study 43 pregnant women from Bosnia and Herzegovina and Croatia were included.

The Project's Common Goals are to Give Answers to

Does and in what capacity the oral health of pregnant women influence pregnancy? Does insufficient oral health of pregnant women (periodontal disease and certain forms of caries) influence gestational age, birth mass of children or oral and cardiovascular health of new-born's, infants and small children? Is chronic inflammation of the oral cavity (periodontal disease and caries) in pregnant women a atherosclerotic and cardiovascular risk factor, that is; do preschool children whose mothers during pregnancy had periodontal disease and/or caries, have a more prominent predictor of early cardiovascular risk (increased body mass index, high value of blood pressure and thickening of intima–media carotids complex) in comparison to children whose mothers during pregnancy had good oral health?

To achieve these goals, we are conducting the research that can last up to 48 months, using a multidisciplinary approach which includes: A gynaecologist, a dentist, a paediatrician, a radiologist, a cardiologist, nutritionists, epidemiologists and statistics. These examinations integrated research from 3 respected centres in Bosnia and Herzegovina and Croatia using combined experience and skills. The aim of this research is to investigate more prominent predictors of early cardiovascular risk increased body mass index, high values of blood pressure and the thickening of the intima-media carotids complex in comparison to children whose mothers had good oral health during pregnancy.

Citation: Dinarević SM, Topić B, Jurišić S, Prohić S, Sporišević L, et al. (2018) The Challenges of Detecting Risk Factors for the Development of Atherosclerosis. J Cardiovasc Dis Diagn 6: 342. doi: 10.4172/2329-9517.1000342

We are presenting the data on the I phase of this Project conducted in: Sarajevo, Mostar, and Split. The plan is to the finish project in 2019 the II phase, and in 2020 the III phase.

The Survey is Designed as a Cohort Study

It included mothers/pregnant women selected by random selection (randomized sampling). During regular gynaecological and obstetric-examination (being I trimester, if necessary, II and III trimester of gestation) a suggestion was given to pregnant women to do their dental examination in order to assess their oral health status. The general health status of the pregnant women was determined on the basis of an assessment of their medical records. The research did not include: Pregnant women with cardiovascular diseases, diabetes mellitus, kidney disease or any chronic illnesses. The survey testing assesses the habits of pregnant women: eating habits, physical activity, alcohol consumption, drugs and smoking.

II phase: The children would be evaluated as new-borns, preterm infants, new-borns of a desirable body weight and new-born of a small birth weight, they would be followed up to their third or fourth year of life. During the systematic review (the first month of life, the first year, the third and fourth year of life) paediatricians should evaluate: the basic characteristics related to pregnancy and childbirth, analysis of the eating habits of children, anthropometric parameters, determine blood pressure values and while radiologists determine the value of complex intima-media carotid artery a cardiologist by echocardiography would evaluate the hemodynamic status of the respondents. The study would include children of proper health conditions, i.e. children with congenital anomalies or certain chronic illnesses would be excluded from the study. Dentists would judge the status of dental health of pregnant women and children and evaluate the appropriateness of the oral health of pregnant women and children. The research is based on the principles of the Helsinki Declaration from 1975 and its amendments in 2008. In order to implement the principles of ethical and bioethical research consent/approval of the appropriate ethics committees/commissions is required. Voluntary inclusion of pregnant women and children is confirmed by signing an informed consent form.

Materials and Methods

After signing the informed consent form i.e. informing mothers/pregnant women, by research methodology the following tests are conducted I phase: - survey testing - dental examination. Evaluation of general health conditions and life habits of pregnant women, determining the basic core characteristics of oral health protection during pregnancy.

N	**43**
Age (mean)	30.7 ± 5.7
Pregnancy status	
Normal	39 (90.70%)
With complications	4 (9.30%)
Education status	
High school	8 (18.60%)
Senior high school	4 (9.30%)
University education	31 (72.10%)
Diseases during pregnancy	
Without	37 (86.05%)
Rare/Lighter illnesses	4 (9.3%)
With bigger complications	2 (4.65%)
How often do you brush your teeth?	
At least 2 × per day	27 (62.8%)
After each meal	16 (37.2%)
How long it takes to brush your teeth?	
Between 1 and 3 min	28 (65.12%)
Longer than 3 min	14 (32.56%)
At most one minute	1 (2.32%)
When do you brush your teeth?	
After each meal	7 (16.3%)
In the morning	1 (2.3%)
In the evening	3 (6.98%)
In the morning and in the evening	32 (74.42%)
The most common reason for dental visit	
Tooth repair	35%
Regular control	53%
Dental pain	12%
How many times did you visit a dentist last year?	
1 ×	23%
2 ×	46%
Not remembering	12%
Didn't visit a dentist	19%

Table 1: Dental status.

Fruit	
>2 per day	65%
1 × per day	30%
Several times per week	5%
Milk and milk products	
>2 per day	33%
1 × per day	49%
1 × per week	0%
Several times per week	9%
Rarely or never	9%
Vegetables	
1 × per day	58%
>2 per day	23%
1 × per week	5%
Several times per week	14%
Juices	
2+ per day	25%
1 × per day	26%
1 × per week	14%
Several times per week	7%
Rarely or never	28%
Sweets	
1 × per day	28%
1 × per week	5%
2+ per day	23%
Several times per week	28%
Rarely or never	16%
Meat	
2+ per day	25%
1 × per day	26%
1 × per week	14%
Several times per week	14%
Rarely or never	21%
Fish	
2+ per day	5%
1 × per day	9%
1 × per week	46%
Several times per week	14%
Rarely or never	26%

Table 2: Eating habits.

Citation: Dinarević SM, Topić B, Jurišić S, Prohić S, Sporišević L, et al. (2018) The Challenges of Detecting Risk Factors for the Development of Atherosclerosis. J Cardiovasc Dis Diagn 6: 342. doi: 10.4172/2329-9517.1000342

KEP index	12.32 ± 5.729		Plaque index		0.3125 ± 0.405	
	S1	S2	S3	S4	S5	S6
A healthy periodental	42.90%	82.10%	39.30%	53.60%	71.40%	35.70%
Bleeding	42.90%	14.30%	46.40%	35.40%	25%	53.60%
Calculus	7.10%	3.60%	7.15%	7.20%	3.60%	10.70%
Periodontal pocket (4-5 mm)	7.10%		7.15%	3.80%	--	--

Table 3: KEP index.

IDB	|__|__||__|				
Date	|__|__| / |__|__| / 2017.				
Institution:					
City:					
State:					
Telephone:	Mail:				
1. GENERAL DATA					
1.1. Birth year					
1.2. Your educational status	Primary school ☐	High school ☐	Higher school ☐	Faculty ☐	☐ Other:
1.3. Your workings status	Employed ☐		Unemployed ☐		
1.4. According to your estimation, your family's economical status is:	Below the average ☐		Average ☐		Above average ☐
1.5. In addition to this child, you:	You have no other children ☐	I have a younger child / children (specify how much)☐ ___			I have an older child / children (how many) ☐
2. PREGNANCY DATA					
2.1. How does pregnancy go?	Orderly ☐	With complications ☐ (name them:)			
2.2. Diagnosed illnesses during pregnancy?	Without ☐		Rearly ☐		Often ☐
2.3. Do you drink alcohol during pregnancy?	Yes ☐		Sometimes ☐		No ☐
2.4. Do you smoke during pregnancy?	Yes ☐		Sometimes ☐		No ☐
Do you use drugs (medicine) during pregnancy?	Yes ☐			No ☐	
*Please name them:					
3. EVALUATION OF KNOWLEDGE ON ORAL HEALTH (Mark one answer to the questions below!)					
3.1 In your opinion, how often you need to brush your teeth?	At least once per day ☐	At least 2x per day ☐	After every meal ☐	It isn't necessary to brush teeth every day ☐	I don't know ☐
3.2. How long it takes to brush your teeth?	At most 1 min ☐	1-3 minutes ☐	Longer than 3 min ☐	I don't know ☐	
3.3. Is it for thorough cleaning of the teeth necessary to use dental floss?	Yes ☐	No ☐	I don't know ☐		
3.4. Is it necessary that the toothpaste contains fluoride contains fluorine?	Yes ☐	No ☐	I don't know ☐		
EVALUATION OF ATTITUDES ABOUT ORAL HEALTH (Mark one of the questions below)					
4.1. Condition of the tooth and the oral cavity have a negative impact on your overall health?	I agree ☐	I disagree ☐	I'm not sure ☐		
4.2. Regular inspection of the teeth and oral cavity is important for the prevention of dental caries and periodontal disease?	I agree ☐	I disagree ☐	I'm not sure ☐		
4.3. Inappropriate state of your dental health can lead to premature birth or having a baby low birth weight?	I agree ☐	I disagree ☐	I'm not sure ☐		
4.4. Inappropriate state of your dental health can lead to problems with dental health of your child or some other disease?	I agree ☐	I disagree ☐	I'm not sure ☐		
4.5. Regular dental examinations are necessary during pregnancy?	I agree ☐	I disagree ☐	I'm not sure ☐		
4.6. Dental interventions are safe during pregnancy?	I agree ☐	I disagree ☐	I'm not sure ☐		
5. EVALUATION OF PRACTICE ON ORAL HEALTH (Mark one of the questions below!)					
5.1. How often do you brush your teeth?	Only in the morning ☐	Only before sleep at night ☐	In the morning and in the night ☐	After every meal ☐	I don't brush teeth every day ☐

Citation: Dinarević SM, Topić B, Jurišić S, Prohić S, Sporišević L, et al. (2018) The Challenges of Detecting Risk Factors for the Development of Atherosclerosis. J Cardiovasc Dis Diagn 6: 342. doi: 10.4172/2329-9517.1000342

5.2. How long do you brush your teeth?	1 minute ☐	1-3 minutes ☐	>3 minutes ☐		
5.3. Do you use dental floss?	Da ☐	Ne ☐			
5.4. Do you use antibacterial mouthwashes?	Da ☐	Ne ☐			
5.5. Do you use chewing gum?	Da ☐	Ne ☐			
5.6. If you use chewing gum, specify which chewing gum you are using?	Chewing gum with xylitol ☐	Chewing gum with sugar ☐	Chewing gum without sugar ☐	I use any kind of chewing gum ☐	
5.7. How many times did you have a regular visit to the dentist last year?	1x ☐	2x ☐	I did not visit dentist ☐	I can not remember ☐	
5.8. The most common reason to visit a dentist is?	regular inspection of the dental health ☐	because of dental pain ☐	tooth extraction ☐	tooth repair ☐	due to other reasons☐ Please, specify:
5.9. How many daily meals do you have?	1-2 ☐	3 meals ☐	4 meals ☐	5 meals ☐	>5 meals ☐

5.10. How often do you consume the following foods?

Grocery	≥ 2 daily	1x day	Several times per week/weekly	1x per week	Rarely or never
Milk or dairy products					
Fruit					
Vegetables					
Fruit juices					
Non-alcoholic drinks					
Sweets (sugar, sweet)					
(Cakes, Biscuits and Chocolate)					
Honey, jam and various spreads					
Candies					
Chips, sticks and other snacks					
Meat					
Fish					
Eggs					

5.11. The most common source of information on tooth health and oral cavity I find out?	From dentist ☐	From medias ☐	Internet (web pages) ☐	Friends and society☐
Signature: _____	Thank you very much for your time!			

Figure 1: Questionnaire for mother: The impact of oral health of pregnant women on the cardiovascular health of children.

Techniques of research

Originally created questionnaires for mothers/pregnant women and children (general data, data from personal history, data on eating habits, data on harmful habits, data on physical activities and data on oral health); determination of initial caries, periodontal examination with complete examination of the oral cavity in mothers and children according to WHO recommendations; determination of body mass and height, calculation of body mass index; determination of blood pressure values in children; determination of carotid complex of intima-media artery by 2D Colour Doppler ultrasonography; 2D Colour Doppler echocardiography evaluation of the cardiovascular system; data base creation in MS Access or MS Office; evaluation of variables by statistical programs in consultation with statisticians; design and development of a web based data collection system for research data entry and analysis.

Survey sheets were originally designed, they include questions about their habits in food intake (frequency, quantity and types) and fluids with a Questioner of dental status in mothers (Figures 1-3).

Anonymous questionnaire for gynecologists and dentists

As a dentist, I avoid the application of local anaesthesia with adrenalin to pregnant women during all months of pregnancy? I avoid any kind of X-raying pregnant women? In pregnancy the extraction of a tooth under pain is not recommended? I avoid therapy of gravidity gingivitis in pregnant women in all phases of pregnancy? Trepanation of teeth, the cause of acute dentogen infection, and incisions are not recommended in pregnant women? The Study Workflow: Pregnant women-gynaecologist-dentist- radiologist- cardiologist-epidemiologist.

Research Data

Preliminary study data I phase (2017-18): Mean age of 43 pregnant women is 30.7 ± 5.7 years; 90.7% pregnancy ran properly; complication detected in 9.3%.

During pregnancy: 86.05% mothers had no new disease diagnosed.

Citation: Dinarević SM, Topić B, Jurišić S, Prohić S, Sporišević L, et al. (2018) The Challenges of Detecting Risk Factors for the Development of Atherosclerosis. J Cardiovasc Dis Diagn 6: 342. doi: 10.4172/2329-9517.1000342

ID NUMBER				
HEALTH CARD NUMBER				
DATE OF EXAM				
A. GENERAL RISK PARAMETERS				
1. COURSE OF PREGNANCY	0 - Normal ☐	1 - Maintained ☐		
2. WEIGHT DURING PREGNANCY	0 - Normal ☐	1 - gained weight over 13 kg ☐	2 - Lost weight ☐	
3. VOMITING, AS AN ACCOMPANYING SYMPTOM OF PREGNANCY:	0 - Rarely/Never ☐	1 - Daily/Often ☐		
4. Dietary regimen during pregnancy:				
4a) milk and dairy products:	0 - Everyday - often ☐	1 - Rarely ☐		
4b) meat-fish-eggs:	0 - Everyday - often ☐	1 - Rarely ☐		
4c) fruit - vegetables:	0 - Everyday - often ☐	1 - Rarely ☐		
4d) sweets (**sugar**, honey, sweet, candy, chocolate, cakes, etc.)	0 - Rarely/Never ☐	1 - Often ☐	2 - Daily ☐	
5. MEDICINE TAKEN IN PREGNANCY	0 - Without ☐	1 - Rarely ☐	2 - Frequent ☐	
6. DIAGNOSED DISEASE DURING PREGNANCY:	0 - Without ☐	1 - Rarely/Smaller ☐	2 -Frequent ☐	
7. FLUOR USED IN THE FORM	0 - Drinking water ☐	1 - Pills from 4. month of pregnancy ☐	2 - Occasionally Pills ☐	3 - Not used ☐
8. DELIVERY:	0 - As scheduled ☐	1 - Early ☐	2 - Prematurity ☐	
9. CHILDS WEIGHT AT BIRTH:	0 - Above 2,5 kg ☐	1 - Less than 2,5 kg ☐		
10. MOTHER'S ORAL HEALTH	0 - Without Illness ☐	1 - Repaired ☐	2 - Non repaired ☐	
11. FATHER'S ORAL HEALTH:	0 - Without Illness ☐	1 - Repaired ☐	2 - Non repaired ☐	
Risk zone	Low Risk	0-8 points		
	Middle Risk	9-16 points		
	High Risk	17-23 points		
B. SPECIFIC RISK PARAMETERS:				
Before health-education interventions				
After the health-education intervention				
0 - No plaque 1 - On the edge of the gingival (probe, staining) 2 - In the gingival sulcus or pocket (visible) 3 - Large quantity				Risk Zone: PI < from Low risk PI from to Middle risk PI > from High risk
B4 - Tooth Status (TS)				
Before health-education interventions				
After the health-education intervention				

Citation: Dinarević SM, Topić B, Jurišić S, Prohić S, Sporišević L, et al. (2018) The Challenges of Detecting Risk Factors for the Development of Atherosclerosis. J Cardiovasc Dis Diagn 6: 342. doi: 10.4172/2329-9517.1000342

H - healthy tooth is present C - Caries R - Tooth root P - Plum CT - Cured tooth E - Tooth removed	Risk zone TS < from Low risk TS from to Middle risk TS > from High risk

Figure 2: Questionaries' of dental status in mothers.

IDB	|__|__||__|			
DATE	|__|__| / |__|__| / 201_.			
INSTITUTION:				
City:				
State:				
Telephone:	Mail:			
1. GENERAL DATA				
1.1. Child's birth date				
1.2. Sex:	Male ☐	Female ☐		
2. INFORMATION ABOUT PREGNANCY AND BIRTH (Mark one of the questions below!)				
2.1. Duration of pregnancy in weeks / weeks?				
2.2. Child was born	At time ☐	Earlier ☐	Prematurity ☐	
2.3. Were there any complications during pregnancy?	No ☐	Yes ☐	Name complications:	
2.4. How is delivery done?	Natural ☐	Cesarean section ☐	Forces ☐	Vacuum extraction ☐
2.5. Was there any complications during delivery?	No ☐	Yes ☐	Name complications:	
2.6. APGAR score	1 minute		5 minute	
2.7. Birth weight (BW) of the child		Birth length (BL) of the child		
3. ABOUT INFANT PERIOD (Mark one of the questions below!)				
3.1 Did your child in first year of life had any				
-congenital heart disease?	No ☐	Yes ☐		
-other chronic illnesses?	No ☐	Yes ☐		
* If your answer to the previous question was "Yes" to indicate which illnesses:				
3.2. Did your child have any natural feeding / breastfeeding during the first year of life?	Not breastfeeding or is breastfeeding a few months ☐ Exclusive breastfeeding for six months while continuing breastfeeding and the addition of non-milk foods after 6 months of age (fruits, vegetables, meat ...) ☐ The combination of breast feeding infant formula containing non-milk foods after 4 months of age ☐ Milk formula containing non-milk foods after 4 months ☐			
EVALUATION OF ORAL HEALTH KNOWLEDGE (Give one of the questions below)				
4.1. If the mother has cavities or inappropriate oral health - states can affect the appearance of cavities in children?	Yes ☐	No ☐	I don't know ☐	
4.2. Is it necessary to treat tooth decay in young children?	Yes ☐	No ☐	I don't know ☐	
4.3. When children should had the first visit to the dentist?	Between 6 and 12 months of the child's life ☐ In the 1st child's year ☐ In the 2nd childhood year ☐ In the 3rd child's year ☐ When a child needs to go to the first grade ☐ I do not know / I'm not sure ☐			
5. EVALUATION OF ATTITUDE ABOUT ORAL HEALTH (Mark one of the questions below!)				
5.1. Children with early childhood caries later may have a high probability for the decay of permanent teeth?	Yes ☐	No ☐	I'm not sure ☐	
5.2. Do you frequent and prolonged night meals with milk formula can lead to the child's caries?	Yes ☐	No ☐	I'm not sure ☐	
5.3. Does frequent or prolonged administration of sweetened beverages (tea, juice) can lead to baby's caries?	Yes ☐	No ☐	I'm not sure ☐	

Citation: Dinarević SM, Topić B, Jurišić S, Prohić S, Sporišević L, et al. (2018) The Challenges of Detecting Risk Factors for the Development of Atherosclerosis. J Cardiovasc Dis Diagn 6: 342. doi: 10.4172/2329-9517.1000342

5.4. Does the mother's kisses, or tasting food over same spoon can cause children's caries?	Yes ☐	No ☐	I'm not sure ☐	
6. EVALUATION OF PRACTICE ON ORAL HEALTH (Mark one of the questions below!)				
6.1. Have you cleaned gingivas or mouth with piece of gauze swelled in water or paper after the last children's meal?	Daily ☐ Rarely ☐ Never ☐			
6.2. When did you begin brushing your child's teeth with fluoride paste?	Immediately after the eruption of teeth ☐ (month of life:) At the age of (specify in months) ☐ Milk teeth - should not be brushed ☐ I do not know ☐			
6.3. How many times a day do you brush your child's teeth with flourid paste?	Only in the morning ☐ Only in the evening ☐ In the morning and in the vening ☐ Weekly ☐ Don't brush teeth ☐			
6.4. At what age (months of life) did the child have the first visit to the dentist?				Never ☐
6.5. In the previous year, how many times child had dental visits?	1x ☐	2x ☐	None ☐	Multiple ☐
6.6. The most common reason to visit a dentist is?	Regular control ☐	Dental pain ☐	Tooth removal ☐	Tooth repair ☐	Other reasons ☐ Name it:
6.7. How many daily meals do you have?	1-2 ☐	3 ☐	4 broka ☐	5 ☐	>5 ☐
6.8. Specify which liquid usually takes a child?	Milk ☐	Water ☐	Tee ☐	Natural juice ☐	

6.9. How often child consumes following foods?

Food	≥ 2 daily	1x daily	Several times a week / weakly	1x per week	Rearly or never
Milk or dairy products					
Fruit					
Vegetables					
Fruit juice					
Non-alchoholic drinks					
Sweets (sugar,sweet)					
Cakes, Biscuits, Chocolate					
Hney, jam					
Candies					
Chips, sticks and other snakcs					
Meat					
Fish					
Eggs					

6.10. The most common source of information on tooth health and oral cavity I find out?	From a dentist ☐	The media ☐	Internet (web) ☐	Friends, acquaintance ☐
Do you competent pediatrician recommended to take your child to the dentist?				Da ☐ Ne ☐
Signature: _____	Thank you very much!			

Figure 3: Childs questionnaire: The impact of oral health of pregnant women on the cardiovascular health of children.

Only 9.3% of women in pregnancy had rare/lighter illnesses and 4.65% of respondents had bigger complications. Educational status: High school finished 8 (18.60%) mothers, senior high school: 4 (9.30%), university education: 31 (72.10%) pregnant women.

Eating habits: 49% dairy products: daily; fruit 65%: two or more times per day; vegetables 23%: Two or more times per day; meat 51%: daily; with 14% more that eats meat several times per week; fish in 46%: once a week; in 26% rarely or never.

KEP (Cavities/Tooth extraction/seal) index: 12.32 ± 5.7; plaque index 0.312; repaired teeth 65.62%; non-repaired teeth 12.5% (Tables 1-3).

Expected results and their significance

The results of this research will show that a certain number of mothers have an inappropriate oral health status due firstly to insufficient oral hygiene, not adhering to appropriate eating guidelines and insufficient visits to the dentist. Using this research, we are going to show that a certain number of mothers with a bad oral health status deliver preterm new-born's and new-born's with low birth weight. We expect that children whose mothers had a bad oral health status age 3 have a worse oral health status in comparison to children whose mothers had a good oral health status. The group of children who are preterm or born with a low birth weight, age 3/4/years, would have a greater body mass index for their age and sex, greater values of blood pressure and greater thickening of the intima-media complex in comparison to the desired values of these parameters for term new-born's and new-borns of a desirable birth weight, with possible incipient signs of cardiovascular system disease including atherosclerotic, in comparison to the control group.

Citation: Dinarević SM, Topić B, Jurišić S, Prohić S, Sporišević L, et al. (2018) The Challenges of Detecting Risk Factors for the Development of Atherosclerosis. J Cardiovasc Dis Diagn 6: 342. doi: 10.4172/2329-9517.1000342

Result

Regular dental therapy can decrease the frequency of the appearance of caries, periodontal disease in pregnant women, the frequency of prematurity, low birth weight with all its potential complications, decrease the financial costs of neonatal intensive care management and cardiovascular repercussions on a new-born's health.

Discussion and Conclusion

The results so far indicate to the awareness of pregnant women of the importance of oral health and its influence on child development. However, it is necessary to wait until the end of the study to see definitive results, the impact of mother's oral health on the developing foetus. The presented cardiovascular-oral health data base for the Balkan region can be used as a geographic, demographic and epidemiologic source of information for the detection and identification of new potential risk factors of individuals for preterm delivery and possible atherosclerosis development. Primary prevention of atherosclerosis should begin as early as possible, during pregnancy, in childhood, by creating a healthy way of life, which will be able to prevent or at least slow the development of atherosclerosis.

Conflict of Interest

None declared.

References

1. Berenson GS, Blonde CV, Farris RP, Foster TA, Frank GC, et al. (1979) Cardiovascular disease risk factor variables during the first year of life. Am J Dis Child 133: 1049-1057.
2. Berenson GS, Srinivasan SR, Hunter SM, Nicklas TA, Freedman DS, et al. (1989) Risk factors in early life as predictors of adult heart disease: The Bogalusa heart study. Am J Med Sci 298: 141-151.
3. Dinarević S (1994) The pathogenesis of atherosclerosis: A review. Br J Cardiol 67: 241-246.
4. Berenson GS, Srinivasan SR, Bao W, Newman WP, Tracy RE, et al. (1998) Association between multiple cardiovascular risk factors and atherosclerosis in children and young adults. N Engl J Med 338: 1650-1656.
5. Dinarević SM (2017) Prevention of cardiovascular disease from an early age. J Cardiovasc Dis Diagn 5: 2-5.
6. Friedewald VE, Kornman KS, Beck JD, Genco R, Goldfine A, et al. (2009) The American Journal of Cardiology and Journal of Periodontology editors' consensus: Periodontitis and atherosclerotic cardiovascular disease. J Peridontol 80: 1021-1032.
7. Teles R, Wang CY (2011) Mechanisms involved in the association between peridontal diseases and cardiovascular disease. Oral Dis 17: 450-461.
8. Boggess KA, Edelstein BL (2006) Oral health in women during preconception and pregnancy: Implications for birth outcomes and infant oral health. Matern Child Health 10: 169-174.
9. Hemalatha VT, Manigandan T, Sarumathi T, Aarthi Nisha V, Amudhan A (2013) Dental considerations in pregnancy: A critical review on the oral care. J Clin Diagn Res 7: 948.
10. Committee Opinion No. 569 (2013) Oral health care during pregnancy and through the lifespan. American College of Obstetricians and Gynecologists. Obstet Gynecol 122: 417-422.
11. Offenbacher S, Katz V, Fertik G, Collins J, Boyd D, et al. (1996) Periodontal infection as a possible risk factor for preterm low birth weight. J Periodontal 67: 1103-1113.
12. Moore S, Ide M, Coward PY, Randhawa M, Borkowska E, et al. (2004) A prospective study to investigate the relationship between periodontal disease and adverse pregnancy outcome. Br Dent J 197: 251-258.
13. Jeffcoat MK, Geurs NC, Reddy MS, Cliver SP, Goldenberg RL, et al. (2001) Periodontal infection and preterm birth: Results of a prospective study. J Am Dent Assoc 132: 875-880.
14. Silk H, Douglass AB, Douglass JM, Silk L (2008) Oral health during pregnancy. Am Physician 77: 1139-1144.
15. Škrinjarić I (2014) Prevention of oral diseases in children: U: Bralić I. et al. Prevention of pediatric disease, Medical publication, Zagreb pp: 412-431.
16. D. Mardešić, Benjak V Nedonošče (2016) Pediatrics: Schoolbook; In: Mardešić D (ed.) Zagreb pp: 389-397.
17. Crispi F, Bijnens B, Figueras F, Bartrons J, Eixarch E, et al. (2010) Fetal growth restriction results in remodeled and less efficient hearts in children. Circulation 121: 2427-2436.
18. Barker D (2008) Human growth and cardiovascular disease. In the window of opportunity: Pre-pregnancy to 24 months of age. Nestle Nutr Work-shop Ser Pediatr program 61: 21-38.
19. Boggess KA, Urlaub DM, Moos MK, Polinkovsky M, El-Khorazaty J, et al. (2011) Knowledge and beliefs regarding oral health among pregnant women. J Am Dent Assoc 142: 1275-1282.

Paediatr Croat. 2019; 63 (Supl 1): 1-5

Pregled
Review

MOGUĆI NOVI FAKTOR RIZIKA ZA PREMATURITET I KARDIOVASKULARNE BOLESTI U DJECE

SENKA MESIHOVIĆ-DINAREVIĆ, LUTVO SPORIŠEVIĆ, BERISLAV TOPIĆ, VJEKOSLAV KRŽELJ, SANJA JURIŠIĆ, GRIT KIRSTEN-SARIĆ, SENAD SARIĆ, ANES JOGUNČIĆ, SAMIR PROHIĆ, AIDA RAMIĆ*

Parodontna bolest je kronična upalna, imunološki posredovana, bolest visoke učestalosti koja je povezana s aterosklerotskom kardiovaskularnom bolesti i mnogim sistemskim bolestima. Ako se parodontna bolest ne prepozna pravovremeno ili neadekvatno liječi, može dovesti do nastanka aterosklerotskih lezija i kliničke ekspresije aterosklerotske kardiovaskularne bolesti. Nedostatan uvid u moguće patološke implikacije statusa oralnog zdravlja u trudnica za nastanak ateroskleroze kao i prijevremenog poroda, inicirao je međunarodni znanstveno-istraživački projekt, koji se realizira u Bosni i Hercegovini, Republici Hrvatskoj i Njemačkoj, u periodu 2017.-2019./2020. godine, pod nazivom "Utjecaj oralnog zdravlja trudnica na kardiovaskularno zdravlje djece". Svrha projekta je evaluirati povezanost oralnog zdravlja s kardiovaskularnim zdravljem i općim zdravljem, uz osvrt na moguću pojavu prijevremenog poroda i rađanja djece niske porođajne mase. Istraživanje će utvrditi da li je trudnička parodontna bolest mogući novi faktor rizika za preuranjenu kardiovaskularnu bolest u djece. Također se želi ukazati na značaj pravilne higijene usne šupljine i redovitih stomatoloških pregleda trudnica u sprječavanju ili smanjenju pojave trudničke parodontne bolesti i utjecaja na komplikacije trudnoće i mogući preuranjeni kardiovaskularni rizik.

Deskriptori: RIZIK, ATEROSKLEROZA, PREMATURITET

Uvod

Temeljnu ulogu u iniciranju, progresiji i komplikacijama kardiovaskularnih bolesti ima ateroskleroza koja se karakterizira suženjem lumena krvne žile zbog lokalnog zadebljanja unutrašnje stjenke krvne žile uslijed formiranja ateroma (1). Jedan od temeljnih razloga što su bolesti uvjetovane aterosklerozom, i dalje jedan od vodećih uzroka obolijevanja i smrti diljem svijeta, je nedovoljno provođenje preventivnih aktivnosti (2, 3). Primarna prevencija ateroskleroze, koja treba početi još u trudnoći i ranoj dječjoj dobi, uključuje promicanje zdravog stila življenja (zdrava ishrana, izbjegavanje pušenja i izloženosti duhanskom dimu i primjerena tjelesna aktivnost) što će spriječiti ili odložiti nastanak faktora rizika za razvoj aterosklerotske bolesti. Sekundarna prevencija uključuje probir aterosklerotskih faktora rizika što omogućuje pravovremeno otkrivanje djece s povećanim rizikom za razvoj ateroskleroze kako bi se odgovarajućim preventivnim i terapijskim aktivnostima usporila ili odložila aterosklerotska bolest. S obzirom da ne postoji specifični lijek za aterosklerozu, najbolji način sprječavanja ateroskleroze i njenih komplikacija je edukacija i preventivne aktivnosti stanovništva (4).

Osim povezanosti oralnih bolesti sa sistemskim oboljenjima, neki autori ukazuju na moguću poveznicu između parodontne bolesti u trudnica s rizikom prijevremenog poroda i rađanja djeteta niske porođajne mase kao i mogućom preuranjenom kardiovaskularnom bolesti u djece (1, 5-8). Cilj ovoga rada je evaluacija povezanosti oralnog zdravlja s kardiovaskularnom bolesti, prezentiranje utjecaja trudničke parodontne bolesti na mogući prijevremeni porod i rađanje djeteta niske porođajne mase, utjecaja lošeg trudničkog oralnog zdravlja na pojavu faktora rizika za aterosklerozu u djece, kao i prikaza temeljnih aspekata aktualnog međunarodnog znanstveno-istraživačkog projekta pod nazivom "Utjecaj oralnog zdravlja trudnica na kardiovaskularno zdravlje djece".

Povezanost oralnog zdravlja s aterosklerotskom kardiovaskularnom bolesti

Oralne bolesti spadaju među najučestalije kronične nezarazne bolesti tokom cjelokupnog života (9). Oralno zdravlje, ključni je indikator blagostanja i kvalitete života, te je povezano s općim zdravljem (9, 10). Nepravovremeno dijagnosticirana i liječena parodontna bolest prelazi u kroničnu upalnu, imunosno posredovanu bolest karakteriziranu propadanjem parodontnog ligamenta i pripadajuće alveolarne kosti. Najznačajnije parodontopatogene bakterije (Aggregatibacter actinomycetetemcomitans,

*Odbor za kardiovaskularnu patologiju
Odjeljenje medicinskih nauka ANU BiH

Adresa za dopisivanje:
Prof. dr. sc. Senka Mesihović-Dinarević, Akademkinja
Odbor za kardiovaskularnu patologiju
Odjeljenje medicinskih nauka ANU BiH
71000 Sarajevo, Bistrik 7, Bosna i Hercegovina
E-mail: dsenka@bih.net.ba

Porphyromonas gingivilas, Tannarella forsythia, Treponema deneticola i Fusobacterium nucleatum) nagomilavaju se u parodontnom džepu uzrokujući imunosni odgovor koji dovodi do oštećenja potpornih struktura i gubitka zuba (11). Povezanost oralnih bolesti i općeg zdravlja je složena i višestruka, tj. oralne bolesti imaju utjecaja na opće zdravlje, kao što i sistemske bolesti utječu na oralno zdravlje (9, 12, 13). Najveći broj oralnih bolesti dijele zajedničke faktore rizika, uključujući nezdravu prehranu bogatu šećerima, pušenje duhana i prekomjerna konzumacija alkohola, s kardiovaskularnim bolestima, zloćudnim tumorima, kroničnim bolestima dišnog sistema i dijabetes melitusom (9, 10).

Osim navedenih riziko faktora, smatra se da u nastanku parodontne bolesti značaj mogu imati i nasljedni faktori rizika. Znanstvenici smatraju da varijacije DNA sekvenci imaju učinak na individualni rizik za razvoj parodontne bolesti jer su utvrdili da uslijed varijacija genskih sekvenci može biti onemogućena aktivnost alfa defenzina, antimikrobnog peptida prisutnog u neutrofilnim granulocitima, odgovornih za uništenje parodontopatogenih bakterija (14). Munz M. i sur. otkrili su genske lokuse udružene sa sklonosti za agresivnu i kroničnu parodontnu bolest, gdje uslijed poremećaja u prirođenoj i stečenoj imunosti dolazi do parodontne bolesti (14). Najučestalije oralne bolesti, karijes i parodontna bolest, mogu izazvati i pogoršati mnogobrojna sistemska oboljenja kao što su: infektivni endokarditis, miokarditis, koronarna srčana bolest, infarkt miokarda, cerebrovaskularna bolest, pneumonija, kronična opstrukcijska bolest pluća, dijabetes melitus, komplikacija trudnoće (preeklampsija, mrtvorođenost i spontani pobačaj), rađanja djece prije termina i niske porođajne mase, osteoporozu, kronična bubrežna bolest i druga oboljenja (9, 12, 13, 15-17).

Aterosklerotska kardiovaskularna bolest spada među vodeće uzroke obolijevanja, prijevremene smrti i disabiliteta diljem svijeta. Osim nemodificirajućih, modificirajućih i netradicijskih faktora rizika, smatra se da u nastanku ateroskleroze značaj može imati i kronična upala uvjetovana određenim mikroorganizmima (18, 19). Njemački patolog Virchow (1948.), američki znanstvenik Ross (1986.) i švedski znanstvenik Hansson (2009.) smatraju da je ateroskleroza upalna bolest, tj. kronični upalni odgovor stjenke krvne žile na različite forme oštećenja endotela (20, 21). Mnogobrojna istraživanja ukazuju na povezanost parodontne bolesti i aterosklerotske kardiovaskularne bolesti (11, 22, 23). Povišene vrijednosti bioamarkera upale, C-reaktivnog proteina (CRP) faktora tumorske nekroze-alfa (TNF-alfa) i interleukina 6 (IL-6) kod pacijenata s udruženom parodontnom bolesti i aterosklerotskom kardiovaskularnom bolesti, ukazuju da je parodontna bolest mogući faktor rizika za aterosklerotsku kardiovaskularnu bolest (12). Sumarno govoreći, aterosklerotski proces uključuje dislipidemiju, endotelijalnu disfunkciju i permeabilnost i nakupljanje lipoproteina u intimi krvne žile (11).

Pojedinci s predominacijom Aggregatibacter actinomycetetemcomitans, Porphyromonas gingivilas, Tannarella forsythia i Treponema deneticola u zubnom plaku, imaju dva puta veću koncentraciju malih, gustih LDL-čestica (sd-LDL) i apolioproteina B (apoB) u serumu vs. pojedinaca koji nemaju parodontnu bolest (11, 24). Endotelijalna disfunkcija temeljni je faktor u razvoju ateroskleroze, a disfunkcionalan endotel pokazuje protrombogeni, proinflamatorni i proateregeni potencijal. Virulentne parodontopatogene bakterije produciraju lipopolisaharide koji stimuliraju Tool-like receptore (TLR) prisutne na endotelijalnim stanicama, a u daljnjem tijeku aterogeneze dolazi do povećanog nivoa endotelijalnih adhezijskih molekula i TNF-alfa (11). Endotelijalne adhezijske molekule privlače makrofage u subendotelijalni prostor, a zajedno sa TNF-alfa i parodontopatogenim bakterijama djelujući na mehanizme prirođene imunosti, povećavaju permeabilnost endotela (11, 12, 23). Veoma patogena bakterija Aggregatibacter actinomycetetemcomitans uslijed proizvodnje toksičnog proteina leukotoxina (LtxA) može dovesti do endotelijalne apoptoze i povećanog broja endotelijalnih adhezijskih molekula promovirajući endotelijalnu permeabilnost (25).

Određene bakterije u zubnom plaku produciraju protein toplinskog šoka (HSP) koji uvjetuje stvaranje protutijela, te uslijed ukrštene rezistencije s endotelom, dolazi do endotelijalne disfunkcije, po tipu autoimune bolesti i iniciranja aterogeneze (23, 26). Disfunkcionalan endotel postaje permeabilan za lipoproteine i makrofage, trombocite i T- limfocite. Hemotaktični faktori iz endotelijalnih stanica i makrofaga induciraju nakupljanje većeg broja monocita koji fagocitiraju lipoproteine i pretvaraju se u pjenaste stanice, te se formira početna aterosklerotska lezija, masna pruga. Parodontopatogene bakterije zajedno s trombocitnim faktorom rasta (PDGF) i lipoproteinima induciraju migraciju glatkih mišićnih stanica, koje stvaraju gusti ekstracelularni matriks, iz medije u intimu krvne žile (11). Kronično upalno stanje uvjetovano parodontopatogenim bakterijama u interakciji s ostalim faktorima rizika uvjetovat će perzistiranje imunološko-inflamatornih procesa dovodeći do nastanka uznapredovalih aterosklerotskih lezija, tj. nastanka fibrolipidne nakupine (aterom) koji se sastoji od središnje lipidne jezgre koja je prekrivena fibroznom kapom. Ako se javi komplikacija plaka javit će se klinička simptomatologija aterosklerotske kardiovaskularne bolesti. Navedeno ukazuje na značaj preventivnih aktivnosti u očuvanju i unaprjeđenju oralnog zdravlja, a time i kardiovaskularnog i općeg zdravlja stanovništva.

Utjecaj parodontne bolesti na trudnoću

Povišena vrijednost cirkulirajućih gestacijskih hormona tijekom trudnoće uzrokuje visoku prevalenciju gingivitisa i hiperplazije gingive (27). Mnogobrojni autori navode moguću udruženost parodontne bolesti trudnica s rizikom prijevremenog poroda i rađanja djeteta niske porođajne mase (28-31). Smatra se da približno 40% trudnica ima određenu formu parodontne bolesti (16, 32). Trudnice sa parodontnom bolesti pokazuju 2-7 puta veću učestalost prijevremenog poroda (28, 33). Morre i sur. ukazuju da trudnice sa parodontnom bolesti su imale veliki broj fetalnih smrti (34). Istražujući povezanost parodontne bolesti trudnica i prijevremenih poroda Jeffcoat

i sur. utvrdili su da su umjereni ili teški oblik parodontne bolesti u ranoj trudnoći bili udruženi s prijevremenim porodom, neovisno o drugim poznatim faktorima rizika za prijevremeni porod (35).

Virulentne parodontopatogene gram - negativne anaerobne bakterije prisutne u zubnom plaku proizvode lipopolisaharide, djeluju kao endotoksini, kao i drugi toksični produkti koji uzrokuju povišene vrijednosti markera upale poput: interleukina (IL-6, IL-8 i IL-1b), TNF-alfa i prostaglandina E_2 (PGE_2), koji cirkulacijom dospijevaju do maternice i posteljice dovodeći do komplikacija trudnoće (spontani pobačaj i mrtvorođenost) kao i prijevremenog poroda i rađanja djece niske porođajne mase (32, 35, 36). Majke prijevremeno rođene djece ili djece niske porođajne mase imale su značajno povišenu vrijednost PGE_2 u gingivalnoj tekućini vs. majkama terminski rođene djece ili djece poželjne porođajne mase (36).

Osim parodontne bolesti tokom trudnoće može se javiti i veća prevalencija karijesa što je uvjetovano smanjenim pH usne šupljine, izraženijim konzumiranjem rafiniranih šećera i neprimjerene oralne higijene (37). Smatra se da virulentne parodontopatogene gram-negativne anaerobne bakterije, kao i kod parodontne bolesti, mogu produciranjem IL-1b, TNF-alfa i PGE_2 dovesti do prijevremenog poroda i rađanja djece niske porođajne mase kod trudnica s težim oblicima karijesa (32, 35). Uzroci prijevremenog poroda u više od 50% slučajeva nisu poznati, a kao mogući uzroci između ostaloga, navode se socioekonomski faktori, akutna i kronična oboljenja majki, višeplodna trudnoća, porodničarski uzroci, nasljedna oboljenja i placentarni uzroci (38).

Bitno je utvrditi moguće faktore rizika za prijevremeni porod i rađanja djece niske porođajne mase jer je na taj način moguće znatno smanjiti učestalost prijevremenog poroda i rađanja djece niske porođajne mase, čime se smanjuje stopa perinatalne smrtnosti i moguće perinatalne komplikacije. Prijevremeno rođena djeca i novorođenčad niske porođajne mase imaju veću učestalost faktora rizika za kardiovaskularnu bolest vs . terminski rođenoj djeci i djeci poželjne porođajne mase (39). Povišena vrijednost glukokortikoida tijekom intrauterinog perioda uvjetuje u postnatalnom periodu programiranje osovine hipotalamus-hipofiza-nadbubrežna žlijezda što ima temeljnu ulogu u većoj učestalosti faktora rizika za kardiovaskularne bolesti u prijevremeno rođene djece i djece niske porođajne mase vs. terminski rođenoj djeci i djeci poželjne porođajne mase (40).

Smatramo da se u svakodnevnom praktičnom radu nedovoljno provodi probir statusa oralnog zdravlja trudnica. Probirom statusa oralnog zdravlja mogli bi se pravovremeno procijeniti održavanje oralne higijene i identificirati trudnice koje imaju karijes ili parodontnu bolest. Pravovremenim stomatološkim liječenjem može se smanjiti učestalost oralnih bolesti kod trudnica, a u kontekstu gore navedenih razmatranja, mogla bi se smanjiti učestalost prijevremenog poroda i rađanja djece niske porođajne mase. te spriječiti ili smanjiti preuranjeno ispoljavanja faktora za aterosklerotsku kardiovaskularnu bolest.

Znanstveno-Istraživački projekt:
"Utjecaj oralnog zdravlja trudnica na kardiovaskularno zdravlje djece"

S obzirom na to da trudnice i liječnici nedovoljno pažnje posvećuju oralnom zdravlju trudnica, kao i da je nedostatno znanje o utjecaju neprimjerenog oralnog zdravlja trudnica na tijek trudnoće i moguću preuranjenu pojavu rizika za aterosklerotsku kardiovaskualrnu bolest, iniciran je međunarodni znanstveno-istraživački projekt, pod nazivom "Utjecaj oralnog zdravlja trudnica na kardiovaskularno zdravlje djece". Aktualno kohortno istraživanje (prva faza istraživanja završena, druga faza u tijeku) se realizira u Bosni i Hercegovini, Republici Hrvatskoj i Njemačkoj, tijekom 2017.-2019./2020. godine. Između ostalog, ciljevi ovog istraživanja su: utvrditi da li trudnička parodontna bolest utječe na gestacijsku dob i porođajnu masu djece, kardiovaskularno zdravlje dojenčadi i male djece, utvrditi da li je trudnička parodontna bolest faktor rizika za aterosklerotsku kardiovaskularnu bolest, kao i da li predškolska djeca čije su majke imale trudničku parodontnu bolest imaju izraženije prediktore prijevremenog kardiovaskularnog rizika (povećan indeks tjelesne mase, viša vrijednost krvnog tlaka i zadebljanje kompleksa intima-media kariotidnih arterija) vs. djeci čije su majke imale primjereno trudničko oralno zdravlje (4).

U istraživanje su uključene majke i djeca isključivo urednog zdravstvenog stanja. Tokom regularnih pregleda trudnica, prvi odnosno drugi ili treći trimestar trudnoće, stomatološkim pregledom utvrđeno je oralno zdravlje trudnica. Uključujući opći pedijatrijski pregled i stomatološki pregled, skupina djece bit će praćena do njihove treće odnosno četvrte godine života (4). Djeca će biti klasificirana prema njihovoj gestacijskoj starosti i porođajnoj masi, a bit će im određen krvni tlak, ultrazvučnim pregledom procijenjena debljina kompleksa intima-media karotidne arterije, a ehokardiografskim pregledom utvrđen hemodinamski status djece (4). Upitnicima su analizirane životne navike i primjerenost oralne higijene majki i djece.

Smatramo da će istraživanje utvrditi da određeni broj majki uslijed nezdravih životnih navika, nepravilne oralne higijene i nedovoljnih posjeta stomatologu ima parodontnu bolest, kao i da rađaju djecu niže gestacijske dobi i niske porođajne mase sa svim svojim mogućim komplikacijama, što bi smanjilo financijske troškove neonatalne intenzivne njege i terapije kao i kardiovaskularne reperkusije zdravlja novorođenčeta. Određena skupina djece s nižom gestacijskom dobi starosti i niskom porođajnom masom će u dobi od tri ili četiri godine imati veći indeks tjelesne mase, veću vrijednost sistoličkog i dijastoličkog krvnog tlaka i veću debljinu kompleksa intima - media karotidne arterije vs. poželjnim vrijednostima indeksa tjelesne mase, krvnog tlaka i debljine kompleksa intima-media karotidne arterije kod djece rođene u terminu i poželjne porođajne mase.

Preliminarni rezultati Prve faze Projekta (period: 2017.-18.) su: prosječna dob 43 trudnice je 30,7 ± 5,7 godina; 90,7% trudnoća protječe uredno; komplikacije su uočene u 9,3%. Tijekom trudnoće kod 86,05% trudnica nije dijagnosticirana nova bolest. Rijetku/laku bolest imalo je

9,3% trudnica, 4,65% je imalo teže komplikacije. Edukacijski status: završena Visoka Škola 8 (18,60%), Srednja Škola: 4 (9,30%), Sveučilišno obrazovanje: 31 (72,10%) trudnica. Navike u prehrani: 49% majki konzumira mliječne proizvode dnevno; voće 65%: dva ili više puta dnevno; povrće 23%: dva ili više puta dnevno; meso 51%: dnevno; 14% konzumira meso nekoliko puta sedmično; riba 46%: jedan put sedmično; 26% rijetko ili nikada. KEP (Cavities/Tooth extraction/seal) index: 12,32 ± 5,7; plak index 0,312; popravljeni zubi 65,62%; kariozni zubi 12,5%. Ovi rezultati indiciraju svjesnost trudnica o značaju oralnog zdravlja i njegovog utjecaja na razvoj djeteta. Međutim, neophodno je pričekati kraj studije i definitivne rezultate. Prezentirana kardiovaskularna-oralna zdravstvena baza podataka za Balkansku regiju može se primijeniti kao i geografski, demografski i epidemiološki izvor informacija za detekciju i identifikaciju mogućih novih faktora rizika za prematuritet i razvoj ateroskleroze.

Zaključak

Mnogobrojna istraživanja ukazuju da virulentne parodontopatogene bakterije prisutne u zubnom plaku uslijed perzistiranja kroničnog upalnog procesa mogu biti faktor rizika za aterosklerotsku kardiovaskularnu bolest i negativno utjecati na sistemsko zdravlje pojedinca. Trudnička parodontna bolest, nepravovremeno dijagnosticirana i neliječena, kao mogući novi faktor rizika za kardiovaskularnu bolest, može nepovoljno utjecati na tijek trudnoće, tj. trudnoća može rezultirati rađanjem djece prije termina i niske porođajne mase, koja u kasnijoj dobi mogu imati preuranjene prediktore kardiovaskularne bolesti. Primarna prevencija aterosklerotske kardiovaskularne bolesti treba početi još tijekom trudnoće. Adekvatna prenatalna njega treba uključiti i zaštitu trudničkog oralnog zdravlja koja treba biti temeljena na pravilnoj higijeni usne šupljine i redovitim stomatološkim pregledima, što će spriječiti ili smanjiti učestalost pojave trudničke parodontne bolesti i njenog utjecaja na komplikacije trudnoće i mogući preuranjeni kardiovaskularni rizik.

NOVČANA POTPORA/*FUNDING*
Nema/*None*

ETIČKO ODOBRENJE/*ETHICAL APPROVAL*
Nije potrebno/*None*

SUKOB INTERESA/*CONFLICT OF INTEREST*
Autori su popunili *the Unified Competing Interest form na www.icmje.org/coi_disclosure.pdf (dostupno na zahtjev)* obrazac i izjavljuju: nemaju potporu niti jedne organizacije za objavljeni rad; nemaju financijsku potporu niti jedne organizacije koja bi mogla imati interes za objavu ovog rada u posljednje 3 godine; nemaju drugih veza ili aktivnosti koje bi mogle utjecati na objavljeni rad./ *All authors have completed the Unified Competing Interest form at www.icmje.org/coi_disclosure.pdf (available on request from the corresponding author) and declare: no support from any organization for the submitted work; no financial relationships with any organizations that might have an interest in the submitted work in the previous 3 years; no other relationships or activities that could appear to have influenced the submitted work.*

LITERATURA

1. Mesihović-Dinarević S. Prevention of Cardiovascular Disease from an Early Age. J Cardiovasc Dis Diagn. 2017; 5: 268. doi: 10.4172/2329- 9517.1000268
2. McGill HC Jr, McMahan CA. Pathology of atherosclerosis in youth and the cardiovascular risk factors. In: Lauer RM, Burns TL, Daniels SR, editors. Pediatric prevention of atherosclerotic cardiovascular disease. Oxford: University Press; 2006; 3-23.
3. Kavey RE, Daniels SR, Lauer RM, Atkins DL, Hayman LL, Taubert K. American Heart Association guidelines for primary prevention of atherosclerotic cardiovascular disease beginning in childhood. Circulation. 2003; 107 (11): 1562-6.
4. Dinarević SM, Topić B, Jurišić S et al. The Challenges of Detecting Risk Factors for the Development of Atherosclerosis. Journal of Cardiovascular Diseases & Diagnosis, 2018; 6: 6. doi:10.4172/2329-9517.1000342
5. Berenson GS, Blonde CV, Farris RP et al. Cardiovascular Disease Risk Factor Variables During the First Year of Life. Am J Dis Child. 1979; 133 (10): 1049-57. doi:10.1001/archpedi.1979.02130100073015
6. Dinarević S. The pathogenesis of atherosclerosis: A review. Br J Cardiol. 1994; 67: 241-6.
7. Friedewald VE, Kornman KS, Beck JD et al. The American Journal of Cardiology and Journal of Periodontology editors' consensus: Periodontitis and atherosclerotic cardiovascular disease. J Peridontol. 2009; 80: 1021-32.
8. Teles R, Wang CY. Mechanisms involved in the association between peridontal diseases and cardiovascular disease. Oral Dis. 2011; 17: 450-61.
9. World Health Organization. Oral health. Key facts. 2018. (pristupljeno 15.1.2019. Dostupno na : https://www.who.int/news-room/fact-sheets/detail/oral-health
10. NSW Ministry of Health. Oral Health 2020: A Strategic Framework for Dental Health in NSW. 2014. (pristupljeno 15.1.2019.) Dostupno na: https://www.health.nsw.gov.au/oralhealth/Publications/progress-on-oral-health-2020.pdf
11. Bale BF, Doneen AL, Vigerust DJ. High-risk periodontal pathogens contribute to the pathogenesis of atherosclerosis. Postgrad Med J. 2017; 93 (1098): 215-20. doi: 10.1136/postgradmedj-2016-134279.
12. Karami S, Ghobadi N, Pakravan A, Dabirian M, Sobouti F. Periodontal Diseases and Possible Future Cardiovascular Events, Are they Related? An Overview. J. Pediatr. Rev. 2018; 6 (1): 44-8. doi: 10.5812/jpr.11144
13. Dörfer C, Benz C, Aida J, Campard G. The relationship of oral health with general health and NCDs: a brief review. Int Dent J. 2017; 67 (2): 14-8. doi: 10.1111/idj.12360.
14. Munz M, Willenborg C, Richter GM et al. A genome-wide association study identifies nucleotide variants at SIGLEC5 and DEFA1A3 as risk loci for periodontitis. Hum Mol Genet. 2017; 26 (3): 2577-88. doi:10.1093/hmg/ddx151.
15. Friedewald VE, Kornman KS, Beck JD et al. American Journal of Cardiology; Journal of Periodontology. The American Journal of Cardiology and Journal of Periodontology editors' consensus: periodontitis and atherosclerotic cardiovascular disease. J Periodontol. 2009; 80 (7): 1021-32. doi:10.1902/jop.2009.097001
16. Boggess KA, Edelstein BL. Oral health in women during preconception and pregnancy: implications for birth outcomes and infant oral health. Matern Child Health J. 2006; 10 (5): 169-74.
17. Horton AL, Boggess KA. Periodontal disease and preterm birth. Obstet Gynecol Clin North Am. 2012; 39 (1): 17-23. doi:10.1016/j.ogc.2011.12.008.
18. Mitchell RN, Schoen FJ. Atherosclerosis. In: Kumar V, Abbas AK, Fausto N, Aster JC, editors. Robbins & Cotran Pathologic Basis of Disease, 8 izd. Philadelphia: WB Saunders; 2010; 1020-35.
19. Morré SA, Stooker W, Lagrand WK, van den Brule AJ, Niessen HW. Microorganisms in the aetiology of atherosclerosis. J Clin Pathol. 2000; 53 (9): 647-54.
20. Ross R. Atherosclerosis-an inflammatory disease. N Engl J Med. 1999; 340: 115-26.
21. Hansson GK. Atherosclerosis-an immune disease: The Anitschkov Lecture 2007. Atherosclerosis. 2009; 202 (1): 2-10.

22. Beukers NG, van der Heijden GJ, van Wijk AJ, Loos BG. Periodontitis is an independent risk indicator for atherosclerotic cardiovascular diseases among 60 174 participants in a large dental school in the Netherlands. J Epidemiol Community Health. 2017; 71 (1): 37-42. doi:10.1136/jech-2015-206745.

23. Bartova J, Sommerova P, Lyuya-Mi et al. Periodontitis as a risk factor of atherosclerosis. J Immunol Res. 2014; 636893. doi:10.1155/2014/636893

24. Rufail ML, Schenkein HA, Barbour SE, Tew JG, van Antwerpen R. Altered lipoprotein subclass distribution and PAF-AH activity in subjects with generalized aggressive periodontitis. J Lipid Res. 2005; 46 (12): 2752-60. doi:10.1194/jlr.M500389-JLR200

25. Dietmann A, Millonig A, Combes V, Couraud PO, Kachlany SC, Grau GE. Effects of Aggregatibacter actinomycetemcomitans leukotoxin on endothelial cells. Microb Pathog. 2013; 61-62: 43-50. doi:10.1016/j.micpath.2013.05.001.

26. Valtonen VV. Role of infections in atherosclerosis. Am Heart J. 1999; 138 (5 Pt 2): 431-3.

27. Vt H, TM, TS, Nisha VA, AA. Dental considerations in pregnancy - a critical review on the oral care. J Clin Diagn Res. 2013; 7 (5): 948-53.

28. Offenbacher S, Katz V, Fertik G et al. Periodontal infection as a possible risk factor for preterm low birth weight. J Periodontol. 1996; 67 (10): 1103-13.

29. Soroye M, Ayanbadejo P, Savage K, Oluwole A. Association between periodontal disease and pregnancy outcomes. Odontostomatol Trop. 2015; 38 (152): 5-16.

30. Turton M, Africa CWJ. Further evidence for periodontal disease as a risk indicator for adverse pregnancy outcomes. Int Dent J. 2017; 67 (3): 148-56. doi:10.1111/idj.12274.

31. Govindaraju P, Venugopal S, Shivakumar MA, Sethuraman S, Ramaiah SK, Mukundan S. Maternal periodontal disease and preterm birth: A case-control study. J Indian Soc Periodontol. 2015; 19 (5): 512-5. doi:10.4103/0972-124X.164751.

32. American College of Obstretricians and Gynecologists. Oral health care during pregancy and through the lifespan. Committee Opinion No.569. Obstet Gynecol. 2013; 122: 417-22.

33. Reza Karimi M, Hamissi JH, Naeini SR, Karimi M. The Relation Between Maternal Periodontal Status and Preterm and Low Birth Weight Infants in Iran: A Case Control Study. Glob J Health Sci. 2015; 8 (5): 184-8. doi:10.5539/gjhs.v8n5p184.

34. Moore S, Ide M, Coward PY, Randhawa M, Borkowska E, Baylis R, Wilson RF. A prospective study to investigate the relationship between periodontal disease and adverse pregnancy outcome. Br Dent J. 2004; 197 (5): 251-8.

35. Jeffcoat MK, Geurs NC, Reddy MS, Cliver SP, Goldenberg RL, Hauth JC. Periodontal infection and preterm birth: results of a prospective study. J Am Dent Assoc. 2001; 132 (7): 875-80.

36. Offenbacher S, Jared HL, O'Reilly PG et al. Potential pathogenic mechanisms of periodontitis associated pregnancy complications. Ann Periodontol. 199; 3 (1): 233-50. doi:10.1902/annals.1998.3.1.233.

37. Silk H, DouglassAB; Douglass JM, Silk L. Oral health during pregnancy. Am Physician. 2008; 77: 1139-44.

38. Mardešić D, Benjak V. Nedonošče. In: Mardešić D i sur, editors. Pedijatrija. osmo prerađeno i dopunjeno izdanje. Zagreb: Školska knjiga. 2016; 389-90.

39. Crispi F, Bijnens B, Figueras F et al. Fetal growth restriction results in remodeled and less efficient hearts in children. Circulation. 2010; 121 (22): 2427-36. doi:10.1161CIRCULATIONAHA.110.937995.

40. Barker DJ. Human growth and cardiovascular disease. Nestle Nutr Work-shop Ser Pediatr program. 2008; 61: 21-38.

Summary

A NEW POTENTIAL RISK FACTOR FOR PREMATURITY AND CARDIOVASCULAR DISEASES IN CHILDREN

Senka Mesihović-Dinarević, Lutvo Sporišević, Berislav Topić, Vjekoslav Krželj, Sanja Jurišić, Grit Kirsten-Sarić, Senad Sarić, Anes Jogunčić, Samir Prohić, Aida Ramić

Periodontal disease is a chronic inflammatory, immune mediated, high-frequency disease associated with atherosclerotic cardiovascular disease and many systemic diseases. If the periodontal disease is not detected promptly, or inadequately treated, can lead to the formation of atherosclerotic lesions and the clinical expression of atherosclerotic cardiovascular disease. Insufficient insight into the possible pathologic implications of the oral health status in pregnant women for the development of atherosclerosis and premature birth, initiated an international scientific research Project, which is being implemented in Bosnia and Herzegovina, Croatia and Germany in the period 2017-2019/2020, entitled "The impact of oral health of pregnant women on the cardiovascular health of children". The purpose of the Project is to evaluate the relationship of oral health and cardiovascular health and general health, with reference to the possible occurrence of premature birth and the birth of children of low birth weight. The research will determine whether the maternity periodontal disease is a possible new risk factor for premature cardiovascular disease in children. It also wants to highlight the importance of proper oral hygiene and regular dental examinations of pregnant women in preventing or reducing the incidence of maternity periodontal disease and the impact on pregnancy complications and possible premature cardiovascular risk.

Descriptors: RISK, ATHEROSCLEROSIS, PREMATURITY

Primljeno/Received: 22. 2. 2019.
Prihvaćeno/Accepted: 20. 3. 2019.

141st Conference
Scientific Federation

SCIENTIFIC FEDERATION
ABODE FOR RESEARCHERS

3rd International Conference on

Women Health and Breast Cancer

May 30-31, 2019 at Nice, France



Collaborator Partner

Media Partner

EMA European Medical Association

Conference Locate.com | Events Get | Enliven Archive | INICOP | Insight Pharma Reports

Medgadget | Maximum | KC | Tabeeby

Scientific Federation

1-8-506/2/B, Begumpet, Hyderabad-500016, India

T: +91-40-6668 7799 | E: contact@scientificfederation.com

http://scientificfederation.com

SCIENTIFIC FEDERATION
ABODE FOR RESEARCHERS

3rd International Conference on
Women Health and Breast Cancer
May 30-31, 2019 at Nice, France

The Oral Health of Pregnant Women as a New Potential Risk Factor for Prematurity and Cardiovascular Diseases in Children

Senka Mesihovic-Dinarevic*, Lutvo Sporisevic, Berislav Topic, Sanja Jurisic, Senad Saric, Grit Kristen Saric, Vjekoslav Krzelj, Anes Joguncic, Samir Prohic and Aida Ramic
Policlinic Eurofarm, Bosnia and Herzegovina

Pregnancy is a state in which complex physical and physiological changes are present, these have important effects on multiple organ systems. Some authors pointed out the possible connection between periodontal disease in pregnant women with the risk of preterm delivery, newborns of a low gestational age with low birth masses and possible cardiovascular disease. Cardiordiovascular disease atherosclerosis is now one of the leading causes of death in developed countries, it begins in childhood, and goes a long time without manifesting symptoms, increasing with age, it begins to seriously threaten health. Insufficient insight into the possible pathological implications of the oral health status of pregnant women with the premature expression of cardiovascular risk factors in children, initiated this research. The aim of the study is to investigate more prominent predictors of early cardiovascular risk factors: increased body mass index, high values of blood pressure and the thickening of the intima-media carotids complex in comparison to children whose mothers had good oral health during pregnancy. Regular dental therapy/care can decrease the frequency of the occurrence of caries and periodontal disease in pregnant women, the frequency of prematurity and low birth weight with all it's potential complications, decreasing the financial costs of neonatal intensive care management and the cardiovascular repercussions on newborn's health. Primary prevention of atherosclerosis should begin as early as possible, during pregnancy, in childhood, creating a healthy way of life, which will be able to prevent or at least slow the development of atherosclerosis.

Biography


Senka Mesihović-Dinarević paediatric cardiologist: Faculty of Medicine, University of Sarajevo 1982. 1982-2016. Paediatric Clinic. PhD 1991, 1992-1996: Royal Brompton Hospital London. Professor of Paediatrics 2006.1994. AEPC, 1995. BPCA.1995-2011: Lecturer London MRCP; 2014. Director of Discipline for health protection of women and children. Consultant: RCPCH 1997. 2003-2016. Director of Paediatric Clinic Sarajevo.2000 FESC, 2008. Full Member of Academy of Sciences and Arts of Bosnia and Herzegovina, Honorary Doctorate of Letters in Cambridge: 2014. 2016.Member of the European Academy of Science and Arts; 2017. Full Professor University for Peace-United Nations, over 498 papers, cited 218 times in the ICI Web of Science.

EuroSciCon

Senka Mesihovic Dinarevic et al., Arch Med 2019, Volume: 11
DOI: 10.36648/1989-5216-C3-013

Senka Mesihovic Dinarevic, Anes Joguncic Grit Kristen Saric, Liana Cambj Sapunar, Davor Petrovic, Berislav Topic, Vjekoslav Krzelj, Sanja Jurisic Senad Saric, Lutvo Sporisevic, Samir Prohic and Aida Ramic

University of Mostar, Bosnia and Herzegovina



Joint Event on
Nursing Diagnosis, Parasitology & Pediatric Cardiology
December 05-06, 2019, Florence, Italy

THE INFLUENCE OF MOTHER'S NUTRITION AND ORAL HEALTH ON THE CHILD'S BLOOD VESSEL DEVELOPMENT

Introduction: The correlation of oral health and cardiovascular diseases has been evaluated in the past, but not the influence of mother's nutrition, its oral health on cardiovascular foetal system development, which can lead to a slower foetus development, lower birth weight, as well as differences in blood vessels development.

Aim: The aim of our study was to evaluate the relationship between oral health of mothers, dietary habits during pregnancy with the development of the coronary and carotid arteries.

Patients & Methods: The study included 40 pregnant women and their newborns. During pregnancy the dental mother's status and eating habits were analysed. After delivery a detailed colour Doppler echocardiography including carotid vessels has been performed.

Results: There was a significant correlation of LVEDs and mothers eating habits (rho = -0.415, p = 0.044), whereby a better value LVEDs determined in infants whose mothers had worse eating habits. Carotid Intima Media had moderate connection with nutrition intake on both of the carotids, whereby the thicker intima had a pregnant woman with poor eating habits (rho = -0.492, p = 0.03). Fortified is a significant correlation between the diameter of the right coronary artery and the KEP index, where the pregnant woman had better diameters with the smaller blood KEP Index (rho = -0.693, p = 0.047). Diameter of descending aorta had a significant association with the dietary habits, and larger diameters defined in the infants, whose mothers had worse eating habits (rho = -0.508, p = 0.011). Ejection fraction (FC) has a significant correlation with the KEP index values (rho = 0.507, p = 0.014). In mothers with KEP index <10, FS had significant lower values (p = 0,002).

Conclusion: There was a significant connection between diet and oral health status of mothers with some segments of the development of cardiovascular system in infants. It is necessary to extend the study and test inference on a larger sample.

Biography

Senka Mesihovic Dinarevic, Bosnia and Herzegovina: Faculty of Medicine, University of Sarajevo 1982. 1982-2016: Paediatric Clinic-University Clinical Centre /UCC/ Sarajevo. MSc 1985, paediatrician age 30, subspecialty in paediatric cardiology: Sarajevo, Belgrade, London: has completed his PhD from Royal Brompton Hospital London/Imperial College in 1991, 1992-1996.She is a Research fellow and Professor of Paediatrics since 2006 at Faculty of Medicine, University of Sarajevo. 1994: She is a Member of AEPC, 1995: BPCA 1995-2011, Lecturer: London MRCP, 2014: She is the Director of Discipline for health protection of women and children including Ginecology-Obstretics Clinic of UCC Sarajevo. She is a Consultant Paediatric Cardiologist at Royal College of Paediatrics and Child Health (RCPCH) 1997 2003-2016: Director of Paediatric Clinic, UCC Sarajevo.2000 FESC, 2008: She is a Full Member of Academy of Sciences and Arts of Bosnia and Herzegovina, since 2009: She is also the Chairman of the Committee of Cardiovascular pathology Academy of Sciences and Arts of Bosnia and Herzegovina; Honorary Doctorate of Letters in Cambridge, England 2014- 2016. He is also a Member of the European Academy of Science and Arts since 2017. He is a Full Professor at the European Center for Peace and Development of the University for Peace established by the United Nations. He has published over 498 papers, cited 218 times in the ICI Web of Science.

dsenka@bih.net.ba

26th International International Conference on Pediatric Cardiology and Congenital Heart Disease

The influence of mother's nutrition and oral health on the child's blood vessel development


Acc.Prof.dr.med.sci. Senka Mesihović-Dinarević

Florence, Italy, 2nd December 2019.









The influence of mother's nutrition and oral health on the child's blood vessel development

Senka Mesihović-Dinarević*, *Anes Jogunčić Grit Kristen Sarić, Liana Cambj-Sapunar, Davor Petrovic, Berislav Topić, Vjekoslav Krželj, Sanja Jurišić, Senad Saric, Lutvo Sporišević, Samir Prohic, Aida Ramić*

Committee for Cardiovascular Pathology, Academy of Sciences and Arts Sarajevo, Policlinic Eurofarm Sarajevo, Bosnia and Herzegovina

Introduction

The correlation of **oral health and cardiovascular diseases** has been evaluated in the past, but not the influence of mother's nutrition, its oral health on cardiovascular **foetal system development**, which can lead to

- a slower foetus development,
- lower birth weight,
- as well as differences in blood vessels development.

Pregnancy is a state in which complex physical and physiological changes are present, these have important effects on multiple organ systems.

Some authors pointed out the **possible connection** between **periodontal disease in pregnant women** with

- the risk of preterm delivery,
- newborns of a low gestational age with low birth masses and
- possible cardiovascular disease.

The aim of our study

- is to evaluate **the relationship** between
- oral health of mothers,
- dietary habits during pregnancy with
- the development of the coronary and carotid arteries, the cardiovascular system.

Topics:


- Atherosclerosis
- Risk factors
- Oral health
- Pregnancy
- Prematurity
- Research data
- Conclusiones

Topic:

Atherosclerosis

Cardiordiovascular disease Atherosclerosis is now one of **the leading causes of death in developed countries,** it begins in childhood, and goes a long time without manifesting symptoms, increasing with age, it begins to seriously threaten health.

Atherosclerosis /the most frequent disease of the arteries/ is characterized by lumen reduction of blood vessels due to local thickening of internal blood vessels caused by **plaque /atheroma.**



As a cardiovascular disease, atherosclerosis is **an interdisciplinary problem, multifactorial in its etiopathogenesis, course and repercussions,**

demands **the action of:**


- family doctors,
- physicians,
- paediatricians,
- cardiologist,
- paediatric cardiologist,
- neurologist,

- diabetologist,
- endocrinologist,
- cardiac and vascular surgeons,
- neurosurgeon,
- radiologist,
- specialist of nuclear medicine,
- epidemiologist,
- nutricionist,
- pathologist,
- biochemist.

with the aim of **early detection and treatment of atheroscelrosis** as well as the reduction of development of risk factors for coronary artery diseases**.**

- **Pathophysiology** of the disease is to damage cells that line the inner surface of the arteries that endothelium due to chemical or mechanical damage.
- Elevated blood cholesterol, smoking, or elevated homocysteine (now one of the more investigated the causes of atherosclerosis) are examples of **chemical damage** and
- **mechanical damage** to the cells are: **high blood pressure, damage of the catheter during diagnostic procedures or even infections.**

- As **the first stage of development of atherosclerosis** occurs **"fatty streak."**
- It is totally reversible, which means that the cessation of the activities of harmful pathogens, endothelial cells recover completely.
- However, **if exposure to adverse event continues**, atherosclerotic plaque grows and more narrow lumen.
- Consequently, **reducing blood flow and tissue oxygen supply becomes insufficient.**



- **In the blood vessel** itself due to reduced elasticity, an increase of blood pressure occurs.
- **Elevated blood pressure** can lead to: cracking in blood of the court or its stratification with the formation of aneurysm or plaque, which can calcify and slim down the wall of the vessel with the ability of forming a clot.
- For atherosclerotic disease itself **plaque composition** rather than its size is much more important.


- Early precursors of vascular changes – **subclinical atherosclerosis** – warrant special attention as this process can be stabilized or even reversed if treated in time.

- Sonographic Intima Media Thickness measurement of the carotid artery (**cIMT: carotid Intima-Media-Thickness**) is considered **a valid surrogate marker** for **cardiovascular risk** allowing assessment of atherosclerotic changes at a very early stage.



Topic:

Risk factors



- hyperlipidemia,
- hypertension,
- smoking,
- diabetes, high fibrinogen, a male sex at younger and middle age,
- menopause in women taking oral contraceptives or hormone replacement therapy only with presence of other risk factors,
- excessive weight, increased level of homocysteine,
- physical inactivity, heredity and immune response in some diseases.

- **The primary intention of prevention** is to **preclude the occurrence of risk factors** for atherosclerosis.
- **The secondary** is **to prevent the development or aggravation of the illness** along with **the reduction or control of existing risks.**
- **Primary prevention should begin as early as possible**, during pregnancy, in childhood, creating a healthy diet, eliminating smoking, regular physical activity, which will prevent or at least slow the development of atherosclerosis.

The consequences of atherosclerosis are:

- **coronary or ischemic heart disease,**
- **myocardial infarction,**
- **cerebrovascular disease and cerebrovascular accident (80% of all heart attacks and brain due to atherosclerosis),**
- **narrowing or blockage of peripheral arteries, carotid arteries,**
- **particularly the legs, which can lead even to the development of gangrene**

Since
there is no specific cure for atherosclerosis,
the best way to prevent this disease,
is prevention!

Therefore,
it is necessary to establish
a dialogue in cardiovascular medicine!

Topic:

Oral health

The association of oral health with atherosclerotic cardiovascular diseases

- **Oral diseases** are among **the most common chronic non-communicable diseases during the entire life.**
- **Oral health** is **a key indicator of well-being and quality of life,** and is closely connected with **the general health.**

- The non-diagnosed and treated periodontal disease **becomes a chronic inflammatory, immune-mediated disease** characterized by deterioration of the periodontal ligament and associated alveolar bone.
- **Correlation between oral diseases and general health is complex** and **multifaceted.**

The largest number of oral diseases share **common risk factors**, **including unhealthy diet high in sugar, tobacco smoking and excessive alcohol consumption, with cardiovascular disease, malignant tumors, chronic diseases of the respiratory system and diabetes mellitus**.

The oral cavity is an integral part of the human organism and therefore there exists a great connection between oral health and systemic health. Not only do some systemic diseases such as

- diabetes,
- osteoporosis,
- HIV infection,
- trisomy 21

have a predisposition for periodontitis but an opposite applies.

Susceptibility to certain systemic disease is **higher in patients with periodontitis than in healthy people:**

- chronic periodontitis is a risk factor **for future cardiovascular disease**,
- pregnant women with chronic periodontitis have **more frequent preterm birth** and
- **newborns have a low birth weight**.

- The explanation for the **pathophysiological mechanisms** of paradont focus and systemic disease is associated with **elevated levels of circulating pro-inflammatory cytokines** and **prostaglandins** derived from:

- deseased parodont,
- gram negative bacteria and their endotoxin-like substances,
- that appear from subgingival biofilms immediately entering the bloodstream.



The dominant problems of everyday dental practice are: **caries, periodontal disease, occlusal abnormalities**, the relationship of oral and general health and a holistic approach to the patient.



Caries and periodontal disease are of an **infectious etiology** therefore the prevention of dental caries and periodontal disease means **preventing odontogenic focuses.**



A Periodontal pocket is a risk factor for **the development or worsening of systemic - focal disease,** because **the infection is always present in it;** a pocket flora is various, massive, virulent and penetrates the soft wall of the **pocket**, the pocket is under **constant mechanical stimulation** during **chewing, swallowing and speech**, all of which **favor the penetration of bacteria** into circulation and **the formation of transient bacteremia**.



Interdisciplinary cooperation in the elimination of potential negative effects of periodontal infections **will result in better systemic health.**

Dental caries and periodontal disease are **the most common and significant oral disease,** they can cause and aggravate numerous other disease:

- of the cardiovascular system (infective endocarditis, atherosclerosis, myocarditis and myocardial infarction),
- of the respiratory system (pneumonia, chronic obstructive pulmonary disease, bronchial asthma and pulmonary abscess),
- neurological disorders (cerebral infarction and cerebral abscess),
- diabetes mellitus, rheumatoid arthritis,
- Alzheimer's disease,
- complications of pregnancy (preeclampsia, stillbirth, miscarriage),
- preterm births and low birth weight, osteoporosis.

Topic:

Pregnancy

- **Pregnancy** is a state in which there are complex physical and physiological changes, which have important effects on multiple systems of organs. High levels of circulating **estrogen** during pregnancy are associated with high incidence of gingivitis and gingival hyperplasia or certain forms of periodontal disease. It is believed that approximately **40% of pregnant women** have a certain form of **periodontal disease**.

- Offenbacher et al. (1996) first suggested a possible **link** between **periodontal disease and risks for child delivery of a low gestational age or small birth weight.** Many researchers suggest a possible link between periodontal disease in pregnant women with a risk for preterm delivery, respectivelly, the birth of babies with low birth weight.
- Researches show that in **pregnant women** with periodontal disease there is a **2-7 times higher risk for prematurity**.

- That is why dentists need **to motivate, educate and instruct** pregnant women towards **a higher level of oral hygiene** and to repair all dento-oral lesions in dental therapeutic procedures, especially **periodontal pockets,** thereby reducing the number of premature births.

It is considered that **in the development of periodontal disease a hereditary risk factors** could have the importance.

Scientists believe that **DNA sequence variations** have an effect on the individual risk for the development of periodontal disease because they have found that **due to the variation of gene sequences,** the activity of alpha defensins /antimicrobial peptides present in neutrophil granulocytes/ which is responsible for the destructionof periodontopathogenic bacteria, could be disabled.

Munz et al. have **discovered the genetic loci** associated with **susceptibility for chronic and aggressive parodont disease** where due **to disorders in congenital and acquired immunity** leads to development of periodontal disease.

Topic:

Prematurity

The incidence of births of premature infants and newborns of a small birth weight is between **5-18%**, depending on the geographical area and population characteristics.


The causes of preterm birth in more than **50%** of cases are not known, and as **possible causes**, among other things, are cited:

- a socioeconomic factors,
- acute and chronic diseases of mothers,
- multiple pregnancy,
- obstetrical causes,
- hereditary diseases and
- placental causes.



- It is important **to identify possible risk factors for preterm birth and birth of children of low birth weight** because in that way **is possible to significantly reduce the incidence of premature birth and the birth of children of low birth weight**, which **decrease the rate of perinatal mortality and possible perinatal complications.**

- Preterm children and infants of low birth weight have **a higher incidence of risk factors for cardiovascular disease** vs. term infants and children of desirable birth weight.



It was established that the mothers who had given birth to premature infants or children of a small birth weight had **significantly elevated levels of PGE2 in the gingival fluid** opposed to the mothers who had given birth to children in the dedicated time or children with an appropriate birth weight.

During pregnancy, there is **an increased susceptibility to caries** due to:

- **the increased acidity of the oral cavity,**
- **the increased consumption of refined sugars and**
- **poor oral hygiene**.

- **Caries bacteria in children** are usually transmitted by direct transmission through the mother's saliva.
- **Mothers** with **high titers of Streptococcus mutans** in their saliva are going to substantially transmit the bacteria to their baby – by **vertical transmission,** creating conditions for early childhood caries.

- Of course, **the time and frequency of the transmission of bacteria, the childs preference for the accumulation of bacteria on it's teeth, the composition and flow of the childs saliva, the amount of refined sugar in the babies food**, are all **significant predictors of early children's caries.**

Due to **the immaturity of their organ systems**, premature babies and infants of a small birth weight are **among in vulnerable group of infants - complications due to prematurity are the leading cause of death in children under 5 years of age**.

- **Preterm children** or **infants of a small birth weight** exhibit a **higher incidence of cardiovascular risk factors (obesity, hypertension, dyslipidemia), and type 2 diabetes mellitus**.

- ***Animal and epidemiological studies*** indicate that conditions of **elevated levels of glucocorticoids intrauterine** during life, **programme the** hypothalamus-pituitary-adrenal gland axis that plays a **key role** in **the higher incidence** of cardiovascular risk in premature infants and children of a small birth weight.

- Apart from **the role of microbiome mouth** (microbiome - all microbes, their genome and mutual interaction in a particular environment) as a **risk factor** for premature delivery or the birth of newborn of small birth weight, it is possible that the **microorganisms of the oral cavity** condition chronic inflammation that can represent an **atherosclerotic cardiovascular risk factor**.

- **Adequate prenatal care** should include **oral health care of pregnant women,** ie. for pregnant women, there is a need to point out the **importance of practicing regular oral hygiene and the need for periodic or as many as or as frequently as needed dental checkups. Any dental intervention or dental radiography** is most appropriate to be undergone in a dental practice **upon the completion of organogenesis,** ie. in **II or III trimester of pregnancy.**



A significant number of pregnant women maintain inappropriate oral health.

- Insufficient visits to the dentist,
- inappropriate oral hygiene and
- the consumption of unhealthy food,

all affect the appearance of **caries as well as parodontal disease** in pregnant women.

- **The sufficient screening/screening of oral health status of pregnant women is not carried out in daily work**, so with screening status of oral health in a greater number of pregnant women, including the assessment of oral hygiene, we would be able to timely identify pregnant women who have **dental caries** respectively **periodontal disease.**
- With **timely dental treatment** we could **reduce** the incidence of dental caries and periodontal disease in pregnant women, and **may reduce** the incidence of preterm delivery and the birth of newborns of a small birth weight, an early childhood caries and predictors of early atherosclerotic cardiovascular risk (**increased body mass index, blood pressure and thickening of the carotid intima-media complex**).



In a cohort of children, who are **preterm or have a low birth weight**,

- a certain number of children age 3 have a greater body mass index,
- a higher value of systolic and diastolic blood pressure, as well as
- a thickening of the intima-media complex of the carotid artery with
- incipient signs of cardiovascular system disease.

Insufficient insight into the possible pathological implications of the oral health status of pregnant women, their eating habits with the premature expression of cardiovascular risk factors in children, initiated this research.

Topic:

Research data

PROJECT

- "Cardiovascular Disease and Oral Health - The impact of oral health of pregnant women on the cardiovascular health of children" a **Project** that is run by the Committee of the Cardiovascular Disease Department of Medical Sciences ASA of Bosnia and Herzegovina, during 2017, the first phase of research was completed according to plan /12 months / as well as second phase /2017-2019/.

- In this study **43 pregnant women from Bosnia and Herzegovina and Croatia** were included, with a plan to also include pregnant women from Slovenia.

An assessment of



THE GENERAL HEALTH OF THE PREGNANT WOMAN



THEIR LIFE HABITS AND



THEIR ORAL HEALTH /FILLING OUT A QUESTION FORM FOR THE FUTURE MOTHERS AND THE DENTAL QUESTIONNAIRES FOR DENTISTS/



AS WELL AS FORMING A DATA BASE IN MS ACCESS/MS OFFICE

The Project's common goals are to give answers to:

- Does and in what capacity **the oral health of pregnant women influence pregnancy ?**
- **Does insufficient oral health of pregnant women** (periodontal disease and certain forms of caries) **influence gestational age, birth mass of children or oral and cardiovascular health** of newborn's, infants and small children?
- **Is chronic inflammation of the oral cavity** (periodontal disease and caries) in pregnant women **a atherosclerotic and cardiovascular risk factor**, that is; do preschool children whose mothers during pregnancy had periodontal disease and/or caries, have a more prominent predictor of early cardiovascular risk (increased body mass index, high value of blood pressure and thickening of intima–media carotids complex) in comparison to children whose mothers during pregnancy had good oral health ?

To achieve these goals, we are conducting **the research that can** last up to 48 months, using a multidisciplinary approach which includes:

- a gynaecologist,
- a dentist,
- a paediatrician,
- a radiologist,
- a cardiologist,
- nutritionists,
- epidemiologists and
- statistics

These examinations integrated research from 3 respected centres in B&H and Croatia using combined experience and skills.

The aim of this research is to investigate **more prominent predictors of early cardiovascular risk**

- increased body mass index,
- high values of blood pressure and
- the thickening of the intima-media carotids complex

in **comparison** to children whose mothers had good oral health during pregnancy.

These are the results of **the First and Second phase of the Project** within the Southeastern European region: „Cardiovascular disease and Oral health - the influence of pregnant women's oral health on children's cardiovascular health" conducted in: **Sarajevo, Mostar, Split.**

The plan is to the finish project:

- 2020 III phase.







Respondents

- The survey is designed **as a kohorn study**: include **mothers /pregnant women** selected by random selection (randomized sampling).

- **During regular gynecological and obstetric-examination** (being I trimester, if necessary, II and III trimester of gestation) a **suggestion was given** to pregnant women **to do their dental examination in order** to assess their **oral health status.**

- The **general health status** of the **pregnant women** was detemined on the basis of an assesment of their **medical records**.

The research **did not include:**

- pregnant women with cardiovascular diseases,
- diabetes mellitus,
- kidney disease or any chronic illnesses.

The survey testing assesses the **habits of pregnant women:**

- eating habits,
- physical activity,
- alcohol consumption,
- drugs and smoking.

II phase:

The children were evaluated

- as newborns,
- preterm infants,
- newborns of a desirable body weight and
- newborn of a small birth weight,

they would be followed up to **their third or fourth year of life.**

During the systematic review (the first month of life, the first year, the third and fourth year of life) **pediatricians** evaluate:

- the basic characteristics related to pregnancy and childbirth,
- analysis of the eating habits of children,
- anthropometric parameters,
- determine blood pressure values and
- while radiologists determine the value of complex intima-media carotid artery
- a cardiologist by echocardography evaluate the hemodynamic status of the respondents.

- **The study included children of proper health conditions**, ie. children with congenital anomalies or certain chronic illnesses were excluded from the study.
- **Dentists** judged **the status of dental health of pregnant women and children,** and **evaluate the appropriateness of the oral health of pregnant women and children.**
- The research is based on the principles of the Helsinki Declaration from 1975 and its amendments in 2008.



In order to implement the principles of ethical and bioethical research a consent / approval of **the appropriate ethics committees / commissions** was obtained.



Voluntary inclusion of pregnant women and children is confirmed by **signing an informed consent form.**

Research Methods

After **signing the informed consent form** ie. informing mothers /pregnant women, by research methodology the following tetsts are conducted **/I phase/** :

- survey testing

- dental examination.

In II phase the following examination of **Children** was performed:

- survey testing,


- anthropometric measurements,

- determination of blood pressure,


- dental examination,


- determination of thickness of intima-media complex of the carotid blood vessels by two-dimensional Color Doppler ultrasonography,
- two-dimensional Color Doppler echocardiographic evaluation of the cardiovascular system.












R
Bifurcation
Near wall
ICA
CCA
ECA
Far wall
L
Bifurcation
Near wall
ICA
CCA
ECA
Far wall

Methods


Questionnaires for mothers/pregnant women and children
Dental examination of pregnant women and children
Paediatric examination
Ultrasound of carotid arteries and echocardiography
Cardiological examination
Evaluation of statistical data

Survey sheets were originally designed, they include questions about their habits in food intake (frequency, quantity and types) and fluids, and the frequency and intensity of physical activity.

Questionary for mothers:



UPITNIK

„UTJECAJ ORALNOG ZDRAVLJA TRUDNICA NA KARDIOVASKULARNO ZDRAVLJE DJECE"

Upitnik za majku

ID6	| | | | |
Datum anketiranja	| | | / | | | / 2017.
Zdravstvena ustanova :	
Mjesto/grad	
Država:	
Kontakt telefon:	

1. OPĆ PODACI

1.1. Godina Vašeg rođenja:						
1.2. Vaš obrazovni status	Osnovna škola ☐	Srednja škola ☐	Viša škola ☐	Fakultet ☐	☐Ostalo:	
1.3. Vaš radni status		Zaposlena ☐		Nezaposlena ☐		
1.4. Prema Vašoj procjeni ekonomski status Vaše porodice/obitelji je:		Ispod prosjeka ☐	Prosječan ☐		Iznad prosjeka ☐	
1.5. Osim ovog djeteta, Vi:	Nemate druge djece ☐	Imam mlađe dijete/djecu (navedite koliko) ☐		Imam starije dijete/djecu (navedite koliko) ☐		

2. PODACI O TRUDNOĆI

2.1. Kako protiče trudnoća?	Uredno ☐	Komplikacije ☐ (navedite:)	
2.2. Dijagnosticirana oboljenja tokom trudnoće?	Bez ☐	Bjelačka/astia ☐	Cista/teška ☐
2.3. Da li tokom trudnoće pijete alkohol?	Da ☐	Ponekada ☐	Ne ☐
2.4. Da li tokom turdnoće pušite?	Da ☐	Ponekada ☐	Ne ☐
Da li konzumirate lijekove tokom trudnoće?	Da ☐	Ne ☐	


* Ukoliko ste na pitanje odgovorili sa "Da", navedite koje lijekove uzimate:

3. EVALUACIJA ZNANJA O ORALNOM ZDRAVLJU
(na navedena pitanja navedite jedan odgovor!)

3.1 Prema Vašim mišljenju, kako često je potrebno prati zube?	Najmanje 2x dnevno ☐	Najmanje 1x dnevno ☐	Prije svakog obroka ☐	Nije potrebno svakodnevno prati zube ☐	Ne znam ☐
3.2. Koliko dugo je potrebno prati zube?	Najduže 1 min ☐	1-3 minute ☐	Duže od 3 minute ☐	Ne znam ☐	
3.3. Da li je za temeljito čišćenje zuba potrebno koristiti zubni konac?	Da ☐	Ne ☐	Ne znam ☐		
3.4. Da li je potrebno da pasta kojom perete zube sadrži fluor?	Da ☐	Ne ☐	Ne znam ☐		

EVALUACIJA STAVA O ORALNOM ZDRAVLJU
(Na navedena pitanja navedite jedan odgovor)

4.1. Stanje zuba i usne šupljine može imati negativan utjecaj na Vaše opće zdravlje?	Slažem se ☐	Ne slažem se ☐	Nisam sigurna ☐	
4.2. Redovna kontrola zuba i usne šupljine štiti je radi sprečavanja karijesa i parodontne bolesti?	Slažem se ☐	Ne slažem se ☐	Nisam sigurna ☐	
4.3. Neprimjereno stanje Vašeg zubnog zdravlja može dovesti do prijevremenog poroda ili rađanja djeteta male porođajne mase?	Slažem se ☐	Ne slažem se ☐	Nisam sigurna ☐	



4.4. Neprimjereno stanje Vašeg zubnog zdravlja može dovesti do problema sa zubnim zdravljem kod Vašeg djeteta ili nekih drugih oboljenja?	Slažem se ☐	Ne slažem se ☐	Nisam sigurna ☐
4.5. Redovni stomatološki pregledi su potrebni tokom trudnoće?	Slažem se ☐	Ne slažem se ☐	Nisam sigurna ☐
4.6. Stomatološke intervencije su sigurne tokom trudnoće?	Slažem se ☐	Ne slažem se ☐	Nisam sigurna ☐

5. EVALUACIJA PRAKSE O ORALNOM ZDRAVLJU
(Na navedena pitanja navedite jedan odgovor!)

5.1. Koliko često perete Vaše zube?	Samo ujutro ☐	Samo navečer ☐	Ujutro i navečer ☐	Nakon svakog obroka ☐	Ne perem zube svaki dan ☐
5.2. Koliko dugo perete Vaše zube?	1 minut ☐		1-3 minute ☐	>3 minute ☐	
5.3. Da li osim četkice i paste za zube koristite konac za zube?	Da ☐			Ne ☐	
5.4. Da li koristite antibakterijske vodice za ispiranje usta?	Da ☐			Ne ☐	
5.5. Da li koristite žvakaće gume?	Da ☐			Ne ☐	
5.6. Ako koristite žvakaće gume, navedite koju žvakaću gumu koristite?	Žvakaća guma s ksilotolom ☐	Žvakaća guma sa šećerom ☐	Žvakaća guma bez šećera ☐	Koristim bilo koju žvakaću gumu ☐	
5.7. Koliko ste puta tokom prošle godine imali redovni posjet stomatologu?	1x ☐	2x ☐	Nisam bila u posjeti stomatologu ☐	Ne mogu se sjetiti ☐	
5.8. Najčešći razlog posjeti stomatologu je?	Redovna kontrola zubnog stanja ☐	Zbog zubnih bolova ☐	Vađenje zuba ☐	Popravka zuba ☐	iz drugih razloga☐ Navedite:
5.9. Koliko dnevnih obroka imate?	1-2 ☐	3 obroka ☐	4 obroka ☐	5 obroka ☐	>5 obroka ☐

5.10. Koliko često konzumirate sljedeće namirnice?


Namirnica	≥ 2 dnevno	1x dnevno	Nekoliko puta sedmično/tjedno	1x sedmično/tjedno	Rijetko ili nikada
Mliječni i mliječni proizvodi					
Voće					
Povrće					
Vlaknasti sokovi					
Bezalkoholni napici					
Slatkiši (šećer, slatko)					
Slatkiši (čokolada,keksi i čokolada)					
Med, marmelada i razni namazi					
Kombinovano					
Čips, štapići i ostale grickalice					
Meso					
Riba					
Jaja					

5.11. Najčešći izvor informacija o zdravlju zuba i usne šupljine samnajem?	Od nadležnog stomatologa ☐	Sredstva javnog informisanja (novine, radio, TV) ☐	Internet (web stranice) ☐	Prijatelja, poznanika ☐

Potpis:

Hvala na izdvojenom vremenu!

Questionary of dental status in mothers:



IDENTIFIKACIONI BROJ TRUDNICE:
BROJ ZDRAVSTVENOG KARTONA:
DATUM PREGLEDA:

A. OPŠTI PARAMETRI RIZIKA:

1. FLUOR KORIŠTEN U OBLIKU:	U vodi za piće ☐	Tableta od 4. mjeseca trudnoće ☐	Tablete povremeno ☐	Nije korišten ☐
2. ORALNO ZDRAVLJE MAJKE:	0 - B.O. ☐	1 - Sanirana usta ☐	2 - Nesanirana usta ☐	
3. ORALNO ZDRAVLJE OCA:	0 - B.O. ☐	1 - Sanirana usta ☐	2 - Nesanirana usta ☐	

B. SPECIFIČNI PARAMETRI RIZIKA:

B1 *Procjena stanja oralne higijene*
Plak index (Silness & Loe)

Prije zdravstveno-vasiptne intervencije

16 12 24
44 32 46

Poslije zdravstveno-vasiptne intervencije

16 12 24
44 32 46

0 - Nema plaka
1 - Uz ivicu gingive (detekcija-korištenjem sonde ili bojenjem)
2 - U gingivalnom sulkusu /džepu ili vidljiv na zubu i ivici gingive
3 - Velika količina

B2 – *Procjena stanja parodonta i potrebni tretmani*
CPITN (Community Periodontal Index of Treatment Needs)

Prije zdravstveno-vasiptne intervencije

17 S1 16 S2 11 26 S3 27
47 S6 46 S5 31 36 S4 37

0-zdrav parodont
1-krvarenje (sondiranje <3,5mm)
2- karmenac (sondiranje <3,5mm)
3- parodontalni džep (4-5 mm)
4 – parodontalni džep (>5mm)

0-nije potrebno liječenje
1- instrukcije OH
2- 1+struganje,poliranje korjena
3- 1+struganje,poliranje korjena
4- 1, struganje,poliranje korjena, parodont.kirurgija


Poslije zdravstveno-vasiptne intervencije:

B3 - *Status zuba*
KEP index

K - Karijes
E - Izvađen zub
P - Plombe

KEP =


Childs Questionary:



UPITNIK

„UTJECAJ ORALNOG ZDRAVLJA TRUDNICA NA KARDIOVASKULARNO ZDRAVLJE DJECE"

Upitnik za dijete



5. EVALUACIJA STAVA O ORALNOM ZDRAVLJU
(Na navedena pitanja navedite jedan odgovor!)

6. EVALUACIJA PRAKSE O ORALNOM ZDRAVLJU
(Na navedena pitanja navedite jedan odgovor!)



Pitanja:

Hvala na izdvojenom vremenu!

Anonymous Questionnaire for gynaecologists and dentists:

- As a dentist I avoid the application of local anaesthesia with adrenalin to pregnant women during all months of pregnancy?
- I avoid any kind of X raying pregnant women?
- In pregnancy the extraction of a tooth under pain is not recommended?
- I avoid therapy of gravidity gingivitis in pregnant women in all phases of pregnancy?
- Trepanation of teeth, the cause of acute dentogen infection, and incisions are not recommended in pregnant women?

The Study Workflow:

Pregnant women-gynecologist-dentist-radiologist- cardiologist-epidemiologist

Patients and methods:

The study included 43 pregnant women and their newborns /I phase/

During pregnancy the **dental mother's status and eating habits** were analysed

After delivery a detailed **colour Doppler echocardiography** including **carotid vessels** has been performed

Preliminary study data **I phase** /2017-2018/:


- mean age of 43 pregnant women is 30.7+/-5.7 years,
- 90.3% pregnancy ran properly,
- complication detected in 9,7%

Preliminary study data **II phase** /2018-2019/:
Included: **40 children** due to sufficinet obtained research data


- Echocardiography
- Color Doppler of carotid arteries


Table 1. Demographic characteristics

N	43
AGE	30,7 ±5,7
PREGNANCY STATUS	
NORMAL	39 (90,70%)
WITH COMPLICATIONS	4 (9,30%)
EDUCATION STATUS	
HIGH SCHOOL	8 (19,30%)
SENIOR HIGH SCHOOL	4 (9,70%)
UNIVERSITY EDUCATION	31(71,00%)
DISEASES DURING PREGNANCY	
WITHOUT	37 (87,1%)
RARE/LIGHTER ILNESSES	4 (9,7%)
WITH COMPLICATIONS	2 (3,2%)



• During pregnancy, **87.1%** mothers had no new disease diagnosed.







Eating habits


- dairy products 48.4% : daily,
- fruit 64.5% : two or more times per day,
- vegetables 22.6% : two or more times per day,
- meat 41.9% : daily; meat: in 45.2% few times during week
- fish in 35.2%: once a week; in 35.8% rarely or never


- Only 3.2% of women in pregnancy consumed **Alcohol** and 3.7% of respondents consumed **cigarettes**.


Assessment of oral hygiene


How often do you need to brush your teeth?
At least 2x daily; 27
After each meal; 16


How long it takes to brush your teeth ?
Between 1 and 3 min; 28
Longer than 3 min; 14
At most one minut; 1


How often you brush your teeth ?
7
3
1
32
After each meal
Just in the evening
Just in the morning
In the morning and evening


How many times did you visit a dentist last year?
23%
46%
12%
19%
1x
2x
Not remembering
Without controls at dentist



Dental status

KEP (Cavities/Tooth extraction/seal) index:12.32+/-5.7

plaque index 0.312


repaired teeth 65.62%;

non repaired teeth 12.5%


Table 2. Dental status

KEP Index	12,32±5,729		PLAQUE Index	0,3125±0,405		
	S1	S2	S3	S4	S5	S6
A healthy periodontal	42,9%	82,1%	39,3%	53,6%	71,4%	35,7%
Bleeding	42,9%	14,3%	46,4%	35,7%	25,0%	53,6%
Tartar (Calculus)	7,1%	3,6%	7,1%	7,1%	3,6%	10,7%
Periodontal pocket (4-5 mm)	7,1%		7,1%	3,6%		


Dental status




Dental status




Dental status




Dental status



Dental status



Dental status



Results of the Second phase

Table 3. Demographic characteristics of the newborns

Values are representing frequency with percentage for gender distribution. Months, Body weight on birth, body weight on the examination are represented with median and interquartile range.
From total number of 40 newborn babies 18 of them were males, and 22 of them were females. When we compare their age in months there isn't any significant difference. Body weigh on birth also didn't differ. Body weight on examination was pretty much similar.


Sex	Male N=18 (45%)	Females N=22 (56%)
Months	4,25 (3,50 - 6,0)	4,0 (2,50 - 5,50)
Body weight on birth	3515 (3200 - 3850)	3425 (3170 - 3700)
Body weight on examination	7425 (6580 - 8700)	6730 (6100 - 8170)

There was a significant correlation of **LVEDs** and m**others eating habits** (rho = -0.415, p = 0.044), whereby a better value LVEDs determined in infants whose mothers had worse eating habits.

Carotid Intima Media had moderate connection with nutrition intake on both of the carotids, whereby **the thicker intima** had a pregnant woman with poor eating habits (rho = -0.492, p = 0.03).

Table 4. Demographic characteristics of mothers

Mother characteristics					
Age	25 (22-30)				
Education			Pregnancy status		
Faculty	22	56%	Normal	35	87,50%
High school	16	40%	With complications	3	7,50%
Collage	2	4,00%	No data	2	5,00%
Employment status			Alcohol during pregnancy		
Working	30	75%	Yes	1	3,00%
Unemployed	10	25%	No	39	97,00%
Economic status			Cigarette smoking		
Above average	3	8%	Yes	0	0%
Average	37	92%	Sometimes	5	12%
Other children			No	35	88%
Older and younger child	3	8%			
Younger child	2	4%			
Older child	8	20%			
No other kids	27	68%			


IN MOST CASES (56%) MOTHER HAD **FACULTY EDUCATION**, AND ALSO THEY WERE **WORKING** IN 75% OF CASES.
INCOME WAS AVERAGE IN 92% OF CASES, AND 68% OF THEM DIDN'T HAVE MORE KIDS.
PREGNANCY WAS WITHOUT ANY COMPLICATIONS IN 87,5% CASES.
ONLY 3% OF THEM **USED ALCOHOL** DURING PREGNANCY, BUT 12% OF THEM SOMETIMES WOULD SMOKE **CIGARETTES.**
Values are representing frequency with percentage for gender distribution.
Age is represented in median with interquartile range



Based on **eating habits of mothers** we have divided subjects **in two groups:** bad and good eating habits.

Table 5. Eating habits during pregnancy






Table 6. Eating habits and Echo imaging

Values are shown as median value with interquartile range, p – probability with level of significance set on p<0,05

Variable	Eating habits		
	Bad eating habits Median (IQ range)	Good eating habits Median (IQ range)	p
DACC	4(3,8-4,7)	4(3,5-4,35)	0,323
LACC	4,1(0,03-4,2)	4,1(3,7-4,2)	0,811
CIMT R	0,04(0,03-0,05)	0,03(0,02-0,04)	0,046*
CIMT L	0,04(0,03-0,04)	0,03(0,03-0,03)	0,053
Aortic root	10(9,5-11)	10(8,8-10,7)	0,810
MPA	10(8,2-11,8)	8,5(8-10,5)	0,296
RPA	3,2(2,2-5,8)	2,6(2,3-3,2)	0,411
LPA	2,8(2,3-5,1)	2,8(2,3-2,9)	0,402
RCA	1,4(1,2-1,9)	1,4(1,1-2)	0,509
LCA	1,5(1,3-1,7)	1,4(1,2-1,6)	0,417
Ascending aorta	9,7(9-12)	9(8,8-9)	0,029*
CW Doppler flow over AA	2,475(1,68-4)	1,62(1,38-1,88)	0,300
Descending aorta	7,5(5,8-8,5)	5,8(5-7)	0,021*
CW Doppler Descending aorta	2,335(1,55-7)	1,575(1,37-1,67)	0,171
E wave m/s	1,07(0,9-1,29)	1,2(1,13-1,3)	0,233
A wave m/s	0,82(0,75-0,95)	0,84(0,81-0,93)	0,550
Flow over AP	1,3(1-1,4)	1,48(1,4-1,48)	0,039*
IVSd	4(4-4,4)	4(4-4,3)	0,371
LVEDd	22(19-25)	19(18-20)	0,091
LVEDs	14(12-15,3)	12(11-13)	0,049*
LVPWd	4,1(3,4-4,6)	4,6(4,1-4,6)	0,149
FS%	37(35-39)	38(36-42)	0,489
EF%	69(66-70,9)	69(69-69)	0,983


There is **significant difference in carotid intima media thickness** in **right carotid artery** (p=0,046).


There was an **differences in diameter of descending aorta**, with children whose mothers had bad eating habits **having larger diameters** (p=0,021)
Also **flow over AP** was better in babies from mothers with good eating habits (p=0,039)
Left ventricular end diastolic dimension was significantly bigger in newborns from mothers with bad eating habits (p=0,049)




Table 6. EHO characteristics based on KEP index


Variable	KEP <=10	KEP>10	p
	Median (IQ range)	Median (IQ range)	
DACC	4,3 (3,6-4,6)	3,8 (3,6-4,4)	0,441
LACC	4,2 (3,4-4,2)	4,1 (3,3-4,2)	0,641
CIMT R	0,04 (0,03-0,05)	0,035 (0,03-0,045)	0,495
CIMT L	0,035 (0,03-0,04)	0,035 (0,03-0,04)	0,704
The root of aorta	10,5 (10-11)	9,95 (8,9-10,35)	0,181
MPA	10,5 (8,3-11,8)	8,75 (7,65-11,65)	0,295
RPA	3,2 (2-6,2)	2,6 (2,2-4,6)	0,489
LPA	2,9 (2,3-5,5)	2,45 (2,2-3)	0,274
RCA	1,4 (1,2-2)	1,3 (1,1-1,75)	0,402
LCA	1,5 (1,3-1,8)	1,4 (1,2-1,7)	0,710
Ascending aorta	10,5 (9-11,8)	9,05 (9-10,95)	0,706
CW Doppler flow AA	1,85 (1,68-4)	1,78 (1,5-4)	0,388

Descending aorta	7 (5,7-8,5)	7,5 (5,6-8,2)	1,000
CW Doppler descending aorta	1,67 (1,58-5)	1,57 (1,37-7)	0,412
E wave m/s	1,15 (1,05-1,3)	1,16 (0,9-1,28)	0,558
A wave m/s	0,76 (0,73-0,84)	0,885 (0,815-0,94)	0,130
Flow over AP	1,38 (1,15-1,4)	1,39 (1,17-1,49)	0,422
IVSd	4 (4-4,7)	4,05 (4-4,35)	0,395
LVEDd	20 (19-22)	22,65 (18,5-24,5)	0,620
LVEDs	13 (12-14)	14 (11-14,8)	0,951
LVPWd	4,1 (3,5-4,6)	4,4 (3,95-5)	0,252
FS%	36 (35-37)	39,5 (37,5-42)	0,003
EF%	68,5 (67-69,5)	69,95 (67-71,45)	0,468

During pregnancy a **dental check** up has been performed for mothers, **KEP index** was used as one of the parameters.

Smaller value is better one. Based on that, a value of 10 was used as border one.

Ejection fraction FS% is significantly higher in group of newborns from mothers who had KEP index larger than 10 (p=0,003).


Eating habits classification
Bad eating habits
Good eating habits
FS%
42,5
40,0
37,5
35,0
32,5
30,0
KEP <=10
KEP >10



Eating habits classification
Bad eating habits
Good eating habits
Med FS%
42
40
38
36
42
38,5
36
35,5
KEP <=10
KEP >10
KEPna10

Table 7. The correlation between KEP and Cardiovascular Variables

Correlations		
Variable	Coefficient	KEP index
RCA	rho	-0,693*
	p	0,0047
A wave m/s	rho	,424*
	p	0,044
FS%	rho	,524*
	p	0,01


Only significant correlation are shown

Fortified is a **significant correlation** between the

- **diameter of the right coronary artery and the KEP index,**

- where the pregnant **woman had better diameters with the smaller blood KEP Index** (rho = -0.693, p = 0.047).

As the KEP rise, the values of RCA were worse.


Descending aorta diameter
8,0000
6,0000
4,0000
2,0000
0,0000
7,43
5,94
Diameter of descending aorta had a significant association with the dietary habits, and larger diameters defined in the infants, whose mothers had worse eating habits (rho = -0.508, p = 0.011).
Bad eating habits <8 points
Healthy eating habits 8+ points



Med FS%
40
39
38
37
36
39,50
36,00
Ejection fraction (FS) has a significant correlation with the KEP index values (rho = 0.524, p = 0.014).
In mothers with KEP index <10, FS had significant lower values (p=0,002).
KEP <=10
KEP >10
KEP classification

Table 8. The correlation between Paradontosis and Cardiovascular Echo Variables

Correlations							
Variable	Coefficient	S1	S2	S3	S4	S5	S6
CIMT L	rho	0,331	0,038	0,478	,629**	0,382	,540*
	p	0,195	0,885	0,052	0,007	0,131	0,025
RPA	rho	,496*	0,389	,728**	,749**	,537*	,774**
	p	0,022	0,081	0	0	0,012	0
LPA	rho	,490*	0,323	,747**	,772**	,475*	,761**
	p	0,028	0,165	0	0	0,034	0
RCA	rho	,560**	0,151	,543**	,572**	0,065	,484*
	p	0,007	0,502	0,009	0,005	0,774	0,022
LCA	rho	,552**	0,159	,605**	,629**	0,065	,551**
	p	0,008	0,479	0,003	0,002	0,775	0,008
Ascending aorta	rho	,458*	0,062	,492*	,548**	0,265	,504*
	p	0,032	0,783	0,02	0,008	0,233	0,017
CW Doppler flow on AA	rho	0,355	0,314	,646**	,593**	0,258	,711**
	p	0,148	0,204	0,004	0,009	0,301	0,001
Descending aorta	rho	-0,201	-0,321	-0,068	0,136	-0,036	0,019
	p	0,369	0,146	0,764	0,546	0,873	0,932
CW Doppler descending aorta	rho	0,276	0,164	,587*	,583*	0,242	,620**
	p	0,268	0,517	0,01	0,011	0,334	0,006
Flow over AP	rho	-0,278	-0,264	-,565**	-,556**	-0,219	-,558**
	p	0,21	0,235	0,006	0,007	0,328	0,007
LVEDd	rho	,542**	0,267	,712**	,723**	,456*	,709**
	p	0,009	0,23	0	0	0,033	0
LVEDs	rho	,511*	0,208	,676**	,718**	0,338	,693**
	p	0,015	0,354	0,001	0	0,124	0
LVPWd	rho	-,531*	-0,347	-,698**	-,664**	-0,311	-,698**
	p	0,011	0,114	0	0,001	0,159	0

Conclusions

The presented cardiovascular-oral health data base for the Balkan region can be used as a geographic, demographic and epidemiologic source of information for the detection and identification of new potential risk factors of individuals for preterm delivery and possible atherosclerosis devlopment.

Primary prevention of atherosclerosis should begin as early as possible, during pregnancy, in childhood, by creating a healthy way of life, which will be able to prevent or at least slow the development of atherosclerosis.

The results so far indicate to the awareness of pregnant women of the importance of oral health and its influence on child development.

However, it is necessary to wait until the end of the study to see definitive results, the impact of mothers oral health on the developing fetus.

There was a significant connection between **diet and oral health status of mothers** with **some segments of the development of cardiovascular system** in infants.

It is necessary to extend the study and test inference on a larger sample.

Final results of the research

Regular dental therapy can decrease the frequency of the appearance of caries, peridontal disease in pregnant women, the frequency of prematurity, low birth weight (with all it's potential complications, decrease the financial costs of neonatal intensive care management and cardiovascular repercussions on a newborn's health.


References

1. Pezelj-Ribarić i sur. Oralno zdravlje – uvjet za opće zdravlje. Rijeka; medicinski fakultet Sveučilišta u Rijeci; 2013
2. Teles R, Wang CY. Mechanisms involved in the association between periodontal diseases and cardiovascular disease. Oral Dis 2011;17:450–461.
3. Zoellner H. Dental infection and vascular disease. Semin Thromb Hemost 2011;37:181–192.
4. Kamer AR, Craig RG, Dasanayake AP, Brys M, Glodzik-Sobanska L, de Leon MJ. Inflammation and Alzheimer's disease: possible role of periodontal diseases. Alzheimers Dement 2008;4:242–250.
5. Boggess KA, Edelstein BL. Oral health in women during preconception and pregnancy: implications for birth outcomes and infant oral health. Matern Child Health J 2006; 10 (5 Suppl):S169-174.
6. Hemaltha V, Manigandan T, Sarumathi T, Aaarthi Nisha V, Amudhan A. Dental considerations in pregnancy – a critical review on the oral care . J Clin Diagn Res 2013; 7:938.
7. American College of Obstetricians and Gynecologists. Oral health care during pregnancy and through the lifespan. Committee Opinion No. 569. Obstet Gynecol 2013;122:417–422.
8. Offenbacher S, Katz V, Fertik G, Collins J, Boyd D, Myynor G i sur. Periodontal infection as a possible risk factor of preterm low birth weight . J Periodontal 1996;67:1103-1113.
9. Soroey M, Avanbadejo P, Savage K, Oluwole A. Association between periodontal disease and pregnancy outcomes. Odontostomatol. Trop. 2015; 38 (152):5-16.
10. Turton M, Africa CW. Further evidence for periodontal disease as a risk indicator for adverse pregnancy outcomes. Int Dent J. 2016 Dec 17. doi: 10.1111/idj.12274.
11. Trivedi S, Lal N, Singhal R. Periodontal disaease during pregnancy. J Orofacial Sci 2015; 7:67.
12. Govindaraju P, Venugopal S, Shivakumar MA, Sethuraman S, Ramaiah SK, Mukundan S. Maternal periodontal disease and preterm birth: A case-control study. J Indian Soc Periodontol. 2015;19(5):512-515. doi: 10.4103/0972-124X.164751
13. Reza Karimi M, Hamissi JH, Naeini SR, Karimi M. The Relationship Between Maternal Periodontal Status of and Preterm and Low Birth Weight Infants in Iran: A Case Control Study. Glob J Health Sci. 2015 ;8(5):184-188. doi: 10.5539/gjhs.v8n5p184.
14. Moore S, Ide M, Coward PY, Randhawa M, Borkowska E, Baylis R, Wilson RF. A prospective study to investigate the relationship between periodontal disease and adverse pregnancy outcome. Br Dent J. 2004 Sep 11;197(5):251-258.
15. Ali TB, Abidin KZ. Relationship of periodontal disease to pre-term low birth weight infants in a selected population--a prospective study. Community Dent Health. 2012; 29(1):100-105.
16. Davenport ES, Williams CE, Sterne JA, Murad S, Sivapathasundram V, Curtis MA. Maternal periodontal disease and preterm low birthweight: case-control study. J Dent Res. 2002;81(5):313-318.
17. Offenbacher S, JaredHL, O'Reilly PG, Wells SR, Salvi GE, Lawrence HP et al. Potential pathogenic mechanism of periodontitis pregnancy complications. Ann Perriodontol 1998; 3:233-250.
18. Jeffcoat MK, Guers NC, Reddy MS, Cliver SP, Goldenberg RL, Hauth JC. Periodontal infection and preterm birth: results of a prospective study. J Am Dent Assoc 2001; 132:875-880.

19. Silk H, Douglass AB, Douglass JM, Silk L. Oral health during pregnancy . Am Physician 2008; 77:1139-1144.
20. Haake SK, Newman MG, Nisengard RJ, Sanz M: Periodontal microbiology. Clinical Periodontology. Edited by: Newman MG, Takei HH, Carranza FA. 2002, Philadelphia: Saunders, 96-112.
21. Nair PNR: Pathogenesis of apical periodontitis and the causes of endodontic failures. Crit Rev Oral Biol Med. 2004, 15: 348-381.
22. Škrinjarić I. Prevencija oralnih bolesti u djece: U: Bralić I. i sur. Prevencija bolesti u dječjoj dobi. Medicinska naklada. Zagreb, 2014. str. 412-431.
23. World Health Organization. Preterm birth. Fact sheet. Reviewed November 2016. http://www.who.int/mediacentre/factsheets/fs363/en/ (pristup 10.1.2017.),.
24. Mardešić D., Benjak V. Nedonošče. U: Mardešić D. i sur. Pedijatrija. Školska knjiga; Zagreb, 2016. str. 389-397.
25. Crispi F, Bijnens B, Figueras F, Bartrons J, Eixarch E, Le Noble F et al. Fetal growth restriction results in remodeled and less efficient hearts in children. Circulation 2010; 121 (22): 2427-2436.
26. Barker DJ. Human growth and cardiovascular disease. Nestle Nutr Work-shop Ser Pediatr program 2008; 61:21-38.
27. Mitchell RN, Schoen FJ. Atherosclerosis. U: Kumar V, Abbas AK,Fausto N, Aster JC, (ur.). Robbins & Cotran Pathologic Basis of Disease, 8 izd. Philadephia: WB Saunders; 2010. str. 1020-35.
28. Al-Ghamdi A, Jiman-Fatani AA, El-Banna H. Role of Chlamidia pneumoniae, helicobacter pylori and cytomegalovirus in coronary artery disease. Pak J Pharm Sci 2011; 24(2):95-101.
29. Bearfield C, Davenport ES, Sivapathasandarem V, Allaker RP. Possible association between amniotic fluid micro-organism infection and microflora in the mouth. BJOG. 2002;109:527-533.
30. Dunlop AL, Mulle JG, Ferranti EP, Edwards S, Dunn AB, Corwin EJ. The Maternal Microbiome and Pregnancy Outcomes that Impact Infant Health: A Review. Advances in neonatal care : official journal of the National Association of Neonatal Nurses. 2015;15(6):377-385. doi:10.1097/ANC.0000000000000218.
31. Boggess KA, Urlaub DM, Moos MK, Polinkovsky M, El-Khorazaty J, Lorenz C. Knowledge and beliefs regarding oral health among pregnant women. J Am Dent Assoc 2011; 142(11):1275-1282.
32. Glaziou Philippe. Samsize. Computer sample size and power. Version 06.2003. Dostupno s URL: http://sampsize.sourceforge.net/ (pristup 10.1.2017.).
33. WHO. Child growth standards. WHO Anthro (version 3.2.2., January 2011) http://www.who.int/childgrowth/software/en/
34. World Health Organization. BMI-for - age BOYS . Birth to 5 years (percentiles) Dostupno s URL: http://www.who.int/childgrowth/standards/cht_bfa_boys_p_0_5.pdf?ua=1 (pristup 10.1.2017.).
35. World Health Organization. BMI-for - age GIRLS . Birth to 5 years (percentiles) Dostupno s URL: http://www.who.int/childgrowth/standards/cht_bfa_girls_p_0_5.pdf?ua=1 (pristup 10.1.2017.)
36. Baylor College of medicine. USDA/ARS Children's Nutrition Research Center, Houston, Texas. Age-based Pediatric Blood Pressure Reference Charts . Dostupno s URL: https://www.bcm.edu/bodycomplab/Flashapps/BPVAgeChartpage.html (dostupno 10.1.2017.).

37. National High Blood Pressure Education Program Working Group on High Blood Pressure in Children and Adolescents. The Fourth report on the diagnosis, evaluation and treatment of high blood pressure in children and adolescents. Pediatrics 2004; 114 (2 Suppl 4 th Report) 555-576. (dostupno 10.1.2017.).
38. Rosamond W, Flegal K, Friday G, Furie K, Go A, Greenlund K et al: Heart disease and stroke, statistics-2007 update: a report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. Circulation 2007; 115-e 69-e171.
39. Friedewald VE, Kornman KS, Beck JD, Genco R, Goldfine A, Libby P et al: The American Journal Cardiology and Journal of Peridontology editor's consensus: peridontitis and atherosclerotic cardiovascular disease. J. Peridontol 2009; 80:1021-32.
40. Paraskevas S, Huizinga JD, Loos BG. A systemic review and meta-analyses on C reactive proten in relation to periodontitis. J Clin Periodontol 2008; 35: 277-90.
41. Humphrey Li, Fu R, Buckley DI, Freeman M, helfand M. Peridontal disease and coronary heart disease incidence: a systemic review and meta-analyses. J Gen Inter Med 2008; 23:2079-86.
42. Tonetti MS, D'Aiuto F, Nihali L, Donald A, Storry C, Parkae M et al: Treatment of peridontitis and endothelial function. N Engl J Med 2007: 356:911-20.
43. Pizzo G, La Cara M, Conti Nibali M, Guiglia R. Peridontitis and preterm delivery. A review od the literature. Minerva. Stomatol 2005; 54:1-14.
44. Srinivas SK, Sammel MD, Stamilio DM, Cothier B, Jeffcoat MK; Perry S et al: Peridontal disease and adverse pregnancy outcome: Is there an association? Am J Obstet Gynecol 2009;200:497-e491-98.
45. Meqa et al: Periodontal disease, preterm delivery and preterm low gestational mass. Acta stomatol Croat. 2017;51(1):33-40. DOI: 10.15644/asc51/1/4.
46. Mesihović-Dinarević S:"Prevention of Cardiovascular Disease from an Early Age", Journal of Cardiovascular Disease/Diagnosis, 2017; S:3, vol 5, issue 3, DOI : 10.4172/2329-9517.1000268, p 2-5.
47. Berenson GS, Blonde CV, Faris RP: Cardiovascular disease risk factor variable during the first year of life. American Journal of Disease in Children, 1979; 133:1049-1057.
48. Berenson GS, Srinivasan SR, Mac D et al: Risk factors in early life as predictors of adult heart disease: The Bogalusa heart study, Am.J.Med.Sci.1989; 298:141-151.
49. Geer JC, McGill, He Strong JP: The fine structure of human atherosclerotic lesions, An. J. Pathol. 1961; 38:263-9.
50. Dhuper S, Buddhe S, Patel S. Managing Cardiovascular Risk in Overweight Children and Adolescents. Paediatric Drugs, 2013, 15(3):181-90. doi: 10.1007/s40272-013-0011-y.
51. Berenson GS, Srinivasan SR, Bao, Newman WP, III, Tracy RE, Wattigney WA. Association between multiple cardiovascular risk factors and atherosclerosis in children and young adults. The Bogalusa Heart Study. N Engl J Med 1998; 338:1650–6.
52. Vt H, T M, T S, Nisha V A, A A. Dental considerations in pregnancy-a critical review on the oral care. J Clin Diagn Res. 2013;7(5):948-53.
53. Offenbacher S, Katz V, Fertik G et al. Periodontal infection as a possible risk factor for preterm low birth weight. J Periodontol. 1996 ;67(10 Suppl):1103-13.
54. Soroye M, Ayanbadejo P, Savage K, Oluwole A. Association between periodontal disease and pregnancy outcomes. Odontostomatol Trop. 2015 ;38(152):5-16.
55. Turton M, Africa CWJ. Further evidence for periodontal disease as a risk indicator for adverse pregnancy outcomes. Int Dent J. 2017; 67(3):148-56. doi: 10.1111/idj.12274.

56. Govindaraju P, Venugopal S, Shivakumar MA, Sethuraman S, Ramaiah SK, Mukundan S. Maternal periodontal disease and preterm birth: A case-control study. J Indian Soc Periodontol. 2015;19(5):512-5. doi: 10.4103/0972-124X.164751.
57. American College of Obstretricians and Gynecologists. Oral health care during pregancy and through the lifespan. Committee Opinion No.569. Obstet Gynecol. 2013; 122:417-22.
58. Reza Karimi M1, Hamissi JH, Naeini SR, Karimi M. The Relation Between Maternal Periodontal Status and Preterm and Low Birth Weight Infants in Iran: A Case Control Study. Glob J Health Sci. 2015;8(5):184-8. doi: 10.5539/gjhs.v8n5p184.
59. Boggess KA, Edelstein BL. Oral health in women during preconception and pregnancy: implications for birth outcomes and infant oral health. Matern Child Health J. 2006;10 (5 Suppl):S169-74.
60. World Health Organization. Oral health. Key facts. 2018. (pristupljeno 15.1.2019. Dostupno na : https://www.who.int/news-room/fact-sheets/detail/oral-health
61. NSW Ministry of Health. Oral Health 2020: A Strategic Framework for Dental Health in NSW. 2014. (pristupljeno 15.1.2019.) Dostupno na: https://www.health.nsw.gov.au/oralhealth/Publications/progress-on-oral-health-2020.pdf
62. Bale BF, Doneen AL, Vigerust DJ. High-risk periodontal pathogens contribute to the pathogenesis of atherosclerosis. Postgrad Med J. 2017;93(1098):215-20. doi: 10.1136/postgradmedj-2016-134279.
63. Karami S, Ghobadi N, Pakravan A, Dabirian M, Sobouti F. Periodontal Diseases and Possible Future Cardiovascular Events, Are they Related? An Overview. J. Pediatr. Rev. 2018; 6 (1): 44-8. doi: 10.5812/jpr.11144
64. Dörfer C, Benz C, Aida J, Campard G. The relationship of oral health with general health and NCDs: a brief review. Int Dent J. 2017; 67(Suppl 2):14-8. doi: 10.1111/idj.12360.
65. Munz M, Willenborg C, Richter GM et al. A genome-wide association study identifies nucleotide variants at SIGLEC5 and DEFA1A3 as risk loci for periodontitis. Hum Mol Genet. 2017. 26(3): 2577-88. doi: 10.1093/hmg/ddx151.
66. Friedewald VE, Kornman KS, Beck JD et al. American Journal of Cardiology; Journal of Periodontology. The American Journal of Cardiology and Journal of Periodontology editors' consensus: periodontitis and atherosclerotic cardiovascular disease. J Periodontol. 2009 ; 80(7):1021-32. doi: 10.1902/jop.2009.097001
67. Boggess KA, Edelstein BL. Oral health in women during preconception and pregnancy: implications for birth outcomes and infant oral health. Matern Child Health J. 2006;10 (5 Suppl):S169-74.
68. Horton AL, Boggess KA. Periodontal disease and preterm birth. Obstet Gynecol Clin North Am. 2012;39(1):17-23, vii. doi: 10.1016/j.ogc.2011.12.008.
69. Mardešić D. i Benjak V. Nedonošče. In: Mardešić D. i sur., editors. Pedijatrija. osmo prerađeno i dopunjeno izdanje. Zagreb: Školska knjiga.; 2016. str. 389-90.
70. Crispi F, Bijnens B, Figueras F et al. Fetal growth restriction results in remodeled and less efficient hearts in children. Circulation. 2010; 121 (22): 2427-36. doi: 10.1161 CIRCULATION AHA. 110.937995.
71. Atherosclerosis. 2015 Feb;238(2):380-7. doi: 10.1016/j.atherosclerosis.2014.12.029. Epub 2014 Dec 24.Intima media thickness measurement in children: A statement from the Association for European Paediatric Cardiology (AEPC) Working Group on Cardiovascular Prevention endorsed by the Association for European Paediatric Cardiology.


ARCHIVES OF
DISEASE IN
CHILDHOOD
October 2021 Volume 106 Issue 10
Highlights from this issue:
Morphine or hydromorphone: which should be preferred?
Evolving management of paediatric portal hypertension
Perinatal outcomes among births to women with infection during pregnancy
Improved quality of reporting safety data of medication in paediatric randomised controlled trials
Cystic fibrosis transmembrane conductance regulator modulators for cystic fibrosis: a new dawn?
An Official Journal of the
Royal College of Paediatrics and Child Health
adc.bmj.com
RCPCH
Royal College of
Paediatrics and Child Health
Leading the way in Children's Health
BMJ

Abstracts

physical health of children all while describing the various ways that one can work with children along with some of the challenges that the experts face each day along with some of the children's experiences.

23 PEDIATRICIAN AND EDUCATIONAL REHABILITATOR – ASSOCIATES IN THE PROCESS OF OBSERVATION, DIAGNOSIS AND REHABILITATION OF CHILDREN WITH DEVELOPMENTAL DISABILITIES

[1]Marija Presečki Zmajlović*, [1]Presečki Zmajlović, [2]Mara Modrić, [3]Ivan Razum. [1]*Kindergarten Radost, Jastrebarsko, Croatia*; [2]*Elementary School Sesvetska Sopnica, Sesvete, Croatia*; [3]*Health Center Jastrebarsko, Croatia*

10.1136/archdischild-2021-europaediatrics.23

The number of children with disabilities and persons with disabilities is continuously increasing, which is evident from the data of the Croatian Institute of Public Health. The report on persons with disabilities in the Republic of Croatia, May 2019, lists 45,314 children with disabilities. The pediatrician, as a doctor who deals with the health care of children from birth to the end of adolescence and cares about improving the child's health and eliminating risk factors that threaten his normal growth and development, certainly has children with disabilities and developmental difficulties among his patients. An important part of any systematic examination of a child should be an assessment of psychomotor development and monitoring developmental milestones. When it comes to children with disabilities it can also include information obtained from co-specialists. The competencies of pediatricians include the promptly detection of developmental dilays, knowledge of various developmental disabilities, and also perspective of related co-experts within different systems.

In providing support to children with disabilities, educational rehabilitators are important part of the interdisciplinary diagnostic team, the team of early intervention experts and professional teams in kindergartens and schools. They are a profession whose diagnostic, educational and rehabilitation procedures and programs apply to children with disabilities and persons with disabilities, the population with visual impairment, learning difficulties, attention deficit hyperactivity disorder, motor disorders, chronic diseases and multiple disabilities. The actions of both, the pediatrician and the educational rehabilitator, are aimed at the well-being of the child with risk factors, developmental delays or identified developmental difficulties. Mutual cooperation should include the exchange of key information and trust in the assessment, competencies and recommendations.

It is particularly important in case of children with autism spectrum disorder, which is primarily characterized by difficulties in social communication and interaction and limited, repetitive patterns of behavior, interests, and activities. During the periodic systematic examination in the pediatric clinic, the mere presence of the parents and the learned strategies can mask the characteristic difficulties. In some children, symptoms are present at an early stage of development, but often become more apparent when social demands exceed the child's current abilities. They are more pronounced during the child's time in kindergarten or school, which is then noticed by other professional associates – educational rehabilitator.

If the parent is not concerned about the child's development, finds it difficult to accept the possibility of the difficulty or denies it, sometimes is missing the transfer of key informations in sequence ,,educational rehabilitator – parent – pediatrician', which unfortunately slows down the process of diagnosis and intervention. As an example of good practice and cooperation between a pediatrician and an educational rehabilitator for the purpose of objective informing, we provide and recommend a written opinion about the developmental status of the child.

24 PRESENCE OF LANGUAGE DISORDERS IN SCHOOL AGE

Martina Galeković*. *Centar za odgoj i obrazovanje Velika Gorica*

10.1136/archdischild-2021-europaediatrics.24

Language difficulties refer to impaired language acquisition and impaired language processing and can be an integral part or one of multiple difficulties. The most common among them is developmental language disorder, which refers to the present difficulties in the domain of language. Developmental language disorder is diagnosed when a child, despite orderly general cognitive abilities and a stimulating environment, has difficulty learning his mother tongue, and then any other language.

The paper presents the results of the analysis of the presence of developmental speech disorder in users of psychosocial support services in the Zagreb County. Linguistic comprehension was examined and the obtained results were analyzed by a qualitative methodology.

We emphasize the awareness of parents, professionals and all those involved in the upbringing and education of the child as a prerequisite for partnership with students and providing the necessary support to students in working with difficulties in the field of language comprehension.

General Pediatrics

25 CHILD'S BLOOD VESSEL DEVELOPMENT IN RELATION TO MOTHER'S DIETARY EATING HABITS AND ORAL HEALTH

Senka Mesihovic Dinarevic*, Anes Jogunčić, Kristen Sarić, Lutvo Sporišević, Berislav Topić, Vjekoslav Krželj, Liana Cambj-Sapunar, Davor Petrović, Sanja Jurišić, Senad Sarić, Samir Prohić. *Polyclinic Eurofarm*

10.1136/archdischild-2021-europaediatrics.25

Dietary eating habits and oral health of pregnant women have the potential to affect pregnancy outcomes. Some observational studies have indicated a significant association of the periodontal disease with adverse pregnancy outcomes, such as the risk of preterm delivery and differences in blood vessels development.

The aim of the study was to evaluate the relationship between the dietary eating habits of mothers and their oral health during pregnancy with the development of the coronary and carotid arteries including intima media thickness.

By random selection, 40 pregnant women and their newborns formed a cohort – pilot study. The approval of the ethics committee was obtained. During pregnancy eating habits and dental mother's status were analyzed. After delivery, a

Arch Dis Child: first published as 10.1136/archdischild-2021-europaediatrics.25 on 11 October 2021. Downloaded from http://adc.bmj.com/ on October 11, 2021 by guest. Protected by copyright.

detailed color Doppler echocardiography including carotid vessels has been performed Newborns from mothers with bad eating habits had significantly higher values of the right carotid artery intima-media thickness/CAIMT/(p=0.046). Oral health was examined with DMF index/Decayed, Missing, Filled/. A significant correlation between the diameter of the right coronary artery and the DMF index was proven, where the pregnant woman had better diameters with the smaller DMF Index (rho = -0.693, p = 0.047). CAIMT had a moderate connection with nutrition intake on both of the carotids, whereby the thicker intima had a pregnant woman with poor eating habits (rho = -0.492, p = 0.03). There was a statistically significant difference in the diameter of descending aorta, with larger diameters in children whose mothers had bad eating habits (p=0.021). Flow over AP was better in newborns from mothers with good eating habits (p=0.039). Fraction shortening/FS%/was significantly higher in newborns whose mothers had a larger DMF index (p=0.03). There was a significant correlation of LVEDs/left ventricle end-diastolic diameter/and mothers eating habits (rho = -0.415, p = 0.044), whereby a higher value LVEDs determined in infants whose mothers had worse eating habits Dietary eating habits and oral health of pregnant women have a significant connection with some segments of cardiovascular system development. Eating habits have an impact on the diameter of blood vessels and on flow.

Unhealthy dietary plans will most probably lead to bad oral health and the presence of periodontitis, which could contribute to the CAIMT and the development of atherosclerosis. It is necessary to extend the study and test inference on a larger sample.

26 FAMILIAL HYPERCHOLESTEROLEMIA: A RARE CASE OF EARLY DIAGNOSIS

Ekaterina Dubonosova*, Anastasia Lamasova, Elizaveta Leonova, Alina Pankova, Kamilla Efendieva. *Research Institute of Pediatrics and Child Health CCH RAS, Ministry of Science and Higher Education*

10.1136/archdischild-2021-europaediatrics.26

Familial hypercholesterolemia (FH) is a common life-threatening genetic condition that causes high cholesterol and leads to a much higher-than-normal risk of coronary heart disease (CHD). The heterozygous type is found in about 1 out of 300-500 people, the homozygous type is quite rare in 1 out of 1 million people.

Objective To analyze the clinical case of family hypercholesterolemia in sibs.

Patients and Methods In 3 siblings (from triplets) at the age of 7 years randomly detected new-onset hypercholesterolemia 5.97 mmol/L, 5.65 mmol/L, 6.43 mmol/L. In a second study after 14 days, hypercholesterolemia persists (6.11 mmol/L, 5.67 mmol/L and 6.49 mmol/L, respectively). Two siblings (identical) had high cholesterol levels (4.060 mmol/L and 4.413 mmol/L) due to low density lipoprotein (LDL). The third child with the lowest level of hypercholesterolemia had normal LDL and a high level of high density lipoprotein (HDL). No evidence of secondary hypercholesterolemia (diabetes mellitus, chronic renal insufficiency, hypothyroidism, cholestatic hepatitis, iatrogenic illness) was found.

A mother (42 years old), sticks to a strict diet with a reduced fat content, but hypercholesterolemia persists, statins therapy is not conducted, recommendations for examining children have not been received.

Results Plasma LDL cholesterol level of 4.0 mmol/L or higher in follow-up blood test, provided parents with hypercholesterolemia, confirms the FH in two children from triplets.

At the same time, there are no external physical signs of the disease in children (xanthomas, corneal arch, xanthelasma). This, along with relatively low hypercholesterolemia, suggests a prognostically favorable Heterozygous Familial Hypercholesterolemia. Genetic screening for the presence of FH is not required to confirm the diagnosis, but may be useful if the diagnosis is ambiguous.

A strict diet with a reduced fat content was recommended to patients, as well as supervision of a cardiologist and lipid screening. It was decided that at the age of 8-10 years, while maintaining LDL cholesterol>4.0 mmol/l in follow-up blood test would be observed even on the recommended diet, the treatment with low doses of statins wiould be discussed.

Conclusion Despite the prevalence of FH and the availability of effective treatment, FH is rarely diagnosed in children. This emphasizes the importance of lipid screening in childhood and cascading screening of all members of the patient's family for the prevention of CHD.

27 ANTHROPOMETRIC DIMENSIONS OF THE ARAL SEE REGION (KARAKALPAKSTAN, REPUBLIC OF UZBEKISTAN) NATIVES MAY REFLECT NEGATIVE INFLUENCE OF PESTICIDE ENDOCRINE DISRUPTOR CHEMICALS ON THE POSTNATAL ONTOGENESIS

Andrey P Pugovkin*, Valeriy O Yerkudov, Azat T Matchanov, Kenjabek U Rozumbetov, Ruslan K Dauletov, Sanovar P Esemuratova, Sergey A Lytaev. *St.Petersburg State Pediatric Medical University, Department of human physiology*

10.1136/archdischild-2021-europaediatrics.27

The aim of the study was revelation of the specific tendencies of the physique development in subjects born and grown up at various distances from the former Aral Lake disaster (ALD) as a possible factor determining exposure window of organochlorine pesticides (OCPs) during the body growth. ALD was a combined result of essential climate aridization and anthropogenic pollution with the excessive application of OCPs in a droughty agricultural region.

A complex anthropometric study including measurement of body mass and length, the size of extremities, pelvis and of skinfold thickness was carried out on 310 volunteers of both sexes. All of them had been born in 1990 – 1995 before the start of effective measures for liquidation of the ALD consequences. The volunteers were divided into three groups, attached to geographical zones: first – zone of ALD, northern part of the region around the town of Muynak; second zone, relatively safe – central part, around the city of Nukus; third zone, safe – the southern part of the region.

The comparison of anthropometric parameters was statistically measured using Kruskal-Wallis test and Mann-Whitney criterion including Bonferoni correction for multiple comparison.

Males from the first group possessed statistically valuable lower body mass, arm length, size of the chest and major joints, waist perimeter, skinfold thickness in comparison to subjects from other groups. In contrast, females from the

Arch Dis Child: first published as 10.1136/archdischild-2021-europaediatrics.25 on 11 October 2021. Downloaded from http://adc.bmj.com/ on October 11, 2021 by guest. Protected by copyright.

Child's blood vessel development in relation to mother's dietary eating habits and oral health

Senka Mesihović-Dinarević [1], Anes Jogunčić [2], Kristen Sarić [3], Lutvo Sporišević [4], Berislav Topić [5], Vjekoslav Krželj [6], Liana Cambj-Sapunar [7], Davor Petrović [8], Sanja Jurišić [9], Senad Sarić [10], Samir Prohić [11]

[1, 5] Academy of Sciences and Arts of BiH, [2, 4] Public Health of Canton Sarajevo, [3, 9, 10] Polyclinic Arbor Vitae Mostar, [6, 7, 8] KCU Split Croatia, [11] Dental Clinic Sarajevo

10th Europaediatrics Zagreb, Croatia 7 - 9 October 2021

Introduction

- Dietary eating habits and oral health of pregnant women have the potential to affect pregnancy outcomes. Some observational studies have indicated a significant association of periodontal disease with adverse pregnancy outcomes, such as risk of preterm delivery and differences in blood vessels development.

Methods & Objectives

The aim of the study was evaluate the relationship between the dietary eating habits of mothers and their oral health during pregnancy with the development of the coronary and carotid arteries including intima media thickness.

Patients and methods:
By random selection, 40 pregnant women and their newborns formed a cohort - pilot study. The approval of the ethics committee was obtained. During pregnancy eating habits and dental mother's status were analysed. After delivery a detailed colour Doppler echocardiography including carotid vessels has been performed.

Results

Newborns from mothers with bad eating habits had significantly higher values of the right carotid artery intima-media thickness /CAIMT/ (p=0.046).

Eating habits and Echo imaging

Values are shown as median value with interquartile range, p – probability with level of significance set on p<0,05

Variable	Eating habits		p
	Bad eating habits Median (IQ range)	Good eating habits Median (IQ range)	
DACC	4(3,4-4,7)	4(3,5-4,35)	0,323
LACC	4,1(0,03-4,2)	4,3(3,7-4,2)	0,811
CIMT R	0,040(0,03-0,05)	0,030(0,02-0,04)	0,046*
CIMT L	0,040(0,03-0,04)	0,030(0,03-0,03)	0,053
Aortic root	10(9,5-11)	10(8,8-10,7)	0,810
MPA	10(8,2-11,8)	8,5(8-10,5)	0,396
RPA	3,2(2,2-3,8)	2,6(2,3-3,2)	0,411
LPA	2,8(2,3-3,1)	2,8(2,3-2,9)	0,402
RCA	1,4(1,2-1,9)	1,4(1,1-2)	0,509
LCA	1,5(1,3-1,7)	1,4(1,2-1,6)	0,417
Ascending aorta	9,7(9-12)	9(8,8-9)	0,029*
CW Doppler flow over AA	2,47(1,68-4)	1,62(1,38-1,88)	0,300
Descending aorta	7,5(5,8-8,5)	5,8(5-7)	0,011*
CW Doppler Descending aorta	2,33(1,55-7)	1,57(1,37-1,67)	0,171
E wave m/s	1,07(0,9-1,29)	1,2(1,13-1,3)	0,233
A wave m/s	0,82(0,75-0,95)	0,84(0,81-0,93)	0,550
Flow over AP	1,3(1-1,4)	1,48(1,4-1,48)	0,039*
IVSd	4(4-4,4)	4(4-4,5)	0,371
LVIDd	22(19-25)	19(18-20)	0,093
LVIDs	14(12-15,3)	12(11-13)	0,049*
LVPWd	4,3(3,4-4,6)	4,6(4,1-4,6)	0,149
FS%	37(35-39)	38(36-42)	0,489

The correlation between KEP and Cardiovaskular Variables


Correlations		
Variable	Coefficient	KEP index
RCA	rho	-0,693*
	p	0,0047
A wave m/s	rho	,424*
	p	0,044
FS%	rho	,524*
	p	0,01

Only significant correlation are shown

There was a significant correlation of **LVEDs** and **mothers eating habits** (rho = -0.415, p = 0.044), whereby a better value LVEDs determined in infants whose mothers had worse eating habits.



Carotid Intima Media had moderate connection with nutrition intake on both of the carotids, whereby **the thicker intima** had a pregnant woman with **poor eating habits** (rho = -0.402, p = 0.03).



Conclusions

Dietary eating habits and oral health of pregnant woman have significant connection with some segments of cardiovascular system development. Eating habits have an impact on diameter of blood vessels and on flow. Unhealthy dietary plans will most probably led to bad oral health and presence of periodontitis, which could contribute to the CAIMT and the development of atherosclerosis.

References & Acknowledgements

- Atherosclerosis. 2015 Feb;238(2):380-7. doi: 10.1016/j.atherosclerosis.2014.12.029. Epub 2014 Dec 24. Intima media thickness measurement in children: A statement from the Association for European Paediatric Cardiology (AEPC) Working Group on Cardiovascular Prevention endorsed by the Association for European Paediatric Cardiology.
- Trivedi S, Lal N, Singhal R. Periodontal disaease during pregnancy. J Orofacial Sci 2015; 7:67.
- Mesihović-Dinarević S:"Prevention of Cardiovascular Disease from an Early Age", Journal of Cardiovascular Disease/Diagnosis, 2017; S:3, vol 5, issue 3, DOI : 10.4172/2329-9517.1000268, p 2-5.

FINANCIAL SUPPORT



AbelaPharm

AMICUS
a Swixx BioPharma company



amsal

POLIKLINIKA



ARBOR VITAE
DR. SARIĆ

